


Clinical Practice Guidelines



October 2014 Edition

Emergency Medical Technician

**CLINICAL PRACTICE
GUIDELINES** - 2014 Edition

Practitioner
Emergency Medical Technician

CLINICAL PRACTICE GUIDELINES - 2014 Edition

PHECC Clinical Practice Guidelines

First Edition 2001

Second Edition 2004

Third Edition 2009

Third Edition Version 2 2011

Fourth Edition April 2012

Fifth Edition July 2014

Published by:

Pre-Hospital Emergency Care Council

Abbey Moat House, Abbey Street, Naas, Co Kildare, Ireland

Phone: + 353 (0)45 882042

Fax: + 353 (0)45 882089

Email: info@phecc.ie

Web: www.phecc.ie

ISBN 978-0-9571028-8-0

© Pre-Hospital Emergency Care Council 2014

Permission is hereby granted to redistribute this document, in whole or part, for educational, non-commercial purposes providing that the content is not altered and that the Pre-Hospital Emergency Care Council (PHECC) is appropriately credited for the work. Written permission from PHECC is required for all other uses. Please contact the author: b.power@phecc.ie

CLINICAL PRACTICE GUIDELINES - 2014 Edition

TABLE OF CONTENTS

FOREWORD	4
ACCEPTED ABBREVIATIONS	5
ACKNOWLEDGEMENTS	7
INTRODUCTION	9
IMPLEMENTATION AND USE OF CLINICAL PRACTICE GUIDELINES	10

CLINICAL PRACTICE GUIDELINES

INDEX	12
KEY/CODES EXPLANATION	14
SECTION 1 CARE PRINCIPLES	15
SECTION 2 PATIENT ASSESSMENT	16
SECTION 3 RESPIRATORY EMERGENCIES	21
SECTION 4 MEDICAL EMERGENCIES	25
SECTION 5 OBSTETRIC EMERGENCIES	48
SECTION 6 TRAUMA	50
SECTION 7 PAEDIATRIC EMERGENCIES	59
SECTION 8 PRE-HOSPITAL EMERGENCY CARE OPERATIONS	81
Appendix 1 – Medication Formulary	84
Appendix 2 – Medications & Skills Matrix	100
Appendix 3 – Critical Incident Stress Management	107
Appendix 4 – CPG Updates for Emergency Medical Technicians	109
Appendix 5 – Pre-Hospital Defibrillation Position Paper	115

FOREWORD



The role of the Pre-Hospital Emergency Care Council (PHECC) is to protect the public by independently specifying, reviewing, maintaining and monitoring standards of excellence for the delivery of quality pre-hospital emergency care for people in Ireland. The contents of this clinical publication are fundamental to how we achieve this goal.

Clinical Practice Guidelines have been developed for responders and practitioners to aid them in providing world-class pre-hospital emergency care to people in Ireland.

I would like to thank the members of the Medical Advisory Committee, chaired by Dr Mick Molloy for their efforts and expertise in developing these guidelines. The council acknowledge the work of the PHECC Executive in researching and compiling these Guidelines, in particular Mr Brian Power, Programme Development Officer. I also commend the many responders and practitioners whose ongoing feedback has led to the improvement and creation of many of the Guidelines herein.

The publication of these Guidelines builds on the legacy of previous publications and marks yet another important milestone in the development of care delivered by responders and practitioners throughout Ireland. Despite the difficulties faced by responders and licensed service providers, I am proud that they continue to develop their skills and knowledge to provide safer and more effective patient care.



Mr Tom Mooney, Chair, Pre-Hospital Emergency Care Council

ACCEPTED ABBREVIATIONS

Accepted abbreviations

Advanced Paramedic	AP
Advanced Life Support	ALS
Airway, Breathing & Circulation	ABC
All Terrain Vehicle	ATV
Altered Level of Consciousness	ALoC
Automated External Defibrillator	AED
Bag Valve Mask	BVM
Basic Life Support	BLS
Blood Glucose	BG
Blood Pressure	BP
Basic Tactical Emergency Care	BTEC
Carbon Dioxide	CO ₂
Cardiopulmonary Resuscitation	CPR
Cervical Spine	C-spine
Chronic Obstructive Pulmonary Disease	COPD
Clinical Practice Guideline	CPG
Degree	°
Degrees Centigrade	°C
Dextrose 10% in water	D ₁₀ W
Drop (gutta)	gtt
Electrocardiogram	ECG
Emergency Department	ED
Emergency Medical Technician	EMT
Endotracheal Tube	ETT
Foreign Body Airway Obstruction	FBAO
Fracture	#
General Practitioner	GP
Glasgow Coma Scale	GCS
Gram	g
Milligram	mg
Millilitre	mL

ACCEPTED ABBREVIATIONS

(contd)

Millimole	mmol
Minute	min
Modified Early Warning Score	MEWS
Motor Vehicle Collision	MVC
Myocardial Infarction	MI
Nasopharyngeal airway	NPA
Milliequivalent	mEq
Millimetres of mercury	mmHg
Nebulised	NEB
Negative decadic logarithm of the H ⁺ ion concentration	pH
Orally (per os)	PO
Oropharyngeal airway	OPA
Oxygen	O ₂
Paramedic	P
Peak Expiratory Flow	PEF
Per rectum	PR
Percutaneous Coronary Intervention	PCI
Personal Protective Equipment	PPE
Pulseless Electrical Activity	PEA
Respiration rate	RR
Return of Spontaneous Circulation	ROSC
Revised Trauma Score	RTS
Saturation of arterial oxygen	SpO ₂
ST Elevation Myocardial Infarction	STEMI
Subcutaneous	SC
Sublingual	SL
Systolic Blood Pressure	SBP
Therefore	∴
Total body surface area	TBSA
Ventricular Fibrillation.....	VF
Ventricular Tachycardia.....	VT
When necessary (pro re nata)	prn

ACKNOWLEDGEMENTS

The process of developing CPGs has been long and detailed. The quality of the finished product is due to the painstaking work of many people, who through their expertise and review of the literature, ensured a world-class publication.

PROJECT LEADER & EDITOR

Mr Brian Power, Programme Development Officer, PHECC.

INITIAL CLINICAL REVIEW

Dr Geoff King, Director, PHECC.

Ms Pauline Dempsey, Programme Development Officer, PHECC.

Ms Jacqueline Egan, Programme Development Officer, PHECC.

MEDICAL ADVISORY COMMITTEE

Dr Mick Molloy, (Chair) Consultant in Emergency Medicine

Dr Niamh Collins, (Vice Chair) Consultant in Emergency Medicine, Connolly Hospital Blanchardstown

Prof Gerard Bury, Professor of General Practice, University College Dublin

Dr Seamus Clarke, General Practitioner, representing the Irish College of General Practitioners

Mr Jack Collins, Emergency Medical Technician, Representative from the PHECC register

Prof Stephen Cusack, Consultant in Emergency Medicine, Cork University Hospital

A/Prof Conor Deasy, Consultant in Emergency Medicine, Cork University Hospital, Deputy Medical Director HSE National Ambulance Service

Mr Michael Dineen, Paramedic, Vice Chair of Council

Mr David Hennelly, Advanced Paramedic, Clinical Development Manager, National Ambulance Service

Mr Macartan Hughes, Advanced Paramedic, Head of Education & Competency Assurance, HSE National Ambulance Service

Mr David Irwin, Advanced Paramedic, representative from the Irish College of Paramedics

Mr Thomas Keane, Paramedic, Member of Council

Mr Shane Knox, Education Manager, National Ambulance Service College

Col Gerard Kerr, Director, the Defence Forces Medical Corps

Mr Declan Lonergan, Advanced Paramedic, Education & Competency Assurance Manager, HSE National Ambulance Service

Mr Seamus McAllister, Divisional Training Officer, Northern Ireland Ambulance Service

Dr David McManus, Medical Director, Northern Ireland Ambulance Service

Dr David Menzies, Consultant in Emergency Medicine, Clinical Lead, Emergency Medical Science, University College Dublin

Mr Shane Mooney, Advanced Paramedic, Chair of Quality and Safety Committee

Mr Joseph Mooney, Emergency Medical Technician, Representative from the PHECC register

Mr David O'Connor, Advanced Paramedic, representative from the PHECC register

Dr Peter O'Connor, Consultant in Emergency Medicine, Medical Advisor Dublin Fire Brigade

Mr Cathal O'Donnell, Consultant in Emergency Medicine, Medical Director, HSE National Ambulance Service

Mr Kenneth O'Dwyer, Advanced Paramedic, representative from the PHECC register

Mr Martin O'Reilly, Advanced Paramedic, District Officer Dublin Fire Brigade

Mr Rory Prevett, Paramedic, representative from the PHECC register

Dr Neil Reddy, Medical Director, Code Blue

Mr Derek Rooney, Paramedic, representative from the PHECC register

Ms Valerie Small, Advanced Nurse Practitioner, Chair of Education and Standards Committee.

Dr Sean Walsh, Consultant in Paediatric Emergency Medicine, Our Lady's Hospital for Sick Children, Crumlin

ACKNOWLEDGEMENTS

EXTERNAL CONTRIBUTORS

Ms Diane Brady, CNM II, Delivery Suite, Castlebar Hospital.

Mr Ray Brady, Advanced Paramedic

Mr Joseph Browne, Advanced Paramedic

Dr Ronan Collins, Director of Stroke Services, Age Related Health Care, Adelaide Et Meath Hospital, Tallaght.

Mr Denis Daly, Advanced Paramedic

Mr Jonathan Daly, Emergency Medical Technician

Dr Zelig Gaffney Daly, General Practitioner

Prof Kieran Daly, Consultant Cardiologist, University Hospital Galway

Mr Mark Dixon, Advanced Paramedic

Dr Colin Doherty, Neurology Consultant

Mr Michael Donnellan, Advanced Paramedic

Dr John Dowling, General Practitioner, Donegal

Mr Damien Gaumont, Advanced Paramedic

Dr Una Geary, Consultant in Emergency Medicine

Dr David Janes, General Practitioner

Mr Lawrence Kenna, Advanced Paramedic

Mr Paul Lambert, Advanced Paramedic

Dr George Little, Consultant in Emergency Medicine

Mr Christy Lynch, Advanced Paramedic

Dr Pat Manning, Respiratory Consultant

Dr Adrian Murphy, Specialist Register in Emergency Medicine

Dr Regina McQuillan, Palliative Care Consultant, St Francis Hospice, Raheny

Prof. Alf Nickolson, Consultant Paediatrician

Dr Susan O'Connell, Consultant Paediatrician

Mr Paul O'Driscoll, Advanced Paramedic

Ms Helen O'Shaughnessy, Advanced Paramedic

Mr Tom O'Shaughnessy, Advanced Paramedic

Dr Michael Power, Consultant Anaesthetist

Mr Colin Pugh, Paramedic

Mr Kevin Reddington, Advanced Paramedic

Ms Barbara Shinnars, Emergency Medical Technician

Dr Dermott Smith, Consultant Endocrinologist

Dr Alan Watts, Register in Emergency Medicine

Prof Peter Weedle, Adjunct Prof of Clinical Pharmacy, National University of Ireland, Cork.

Mr Brendan Whelan, Advanced Paramedic

SPECIAL THANKS

HSE National Clinical Programme for Acute Coronary Syndrome

HSE National Asthma Programme

HSE National Diabetes Programme

HSE National Clinical Programme for Emergency Medicine

HSE National Clinical Programme for Epilepsy

HSE National Clinical Programme for Paediatrics and Neonatology

A special thanks to all the PHECC team who were involved in this project. In particular Ms Deirdre Borland for her dedication in bringing this project to fruition.

EXTERNAL CLINICAL PROOFREADING

Ms Joanne Fitzroy, EMT

Ms Niamh O'Leary, EMT

INTRODUCTION




Clinical Practice Guidelines for pre-hospital care are under constant review as practices change, new therapies and medications are introduced, and as more pre-hospital clinical pathways are introduced such as Code STEMI and code stroke which are both leading to significant improved outcomes for patients. A measure of how far the process has developed can be gained from comparing the 29 Standard Operating Procedures for pre-hospital care in existence prior to the inception of the Pre-Hospital Emergency Care Council and the now more than 319 guidelines and growing.

The 2014 guidelines include such new developments as the use of intranasal fentanyl for advanced paramedics and harness induced suspension trauma for both practitioners and responders.

Clinical Practice Guidelines recognise that practitioners and responders provide care to the same patients but to different skill levels and utilising additional pharmaceutical interventions depending on the practitioner level.

This edition of the guidelines has introduced some new concepts such as the basic tactical emergency care standard at EFR and EMT level for appropriately employed individuals. As ever feedback on the guidelines from end users or interested parties is always welcomed and may be directed to the Director of PHECC or the Medical Advisory Committee who review each and every one of the guidelines before they are approved by the Council.



Dr Mick Molloy, Chair, Medical Advisory Committee.

Feedback on the CPGs may be given through the centre for Pre-hospital Research www.ul.ie/cpr/forum

IMPLEMENTATION

Clinical Practice Guidelines (CPGs) and the practitioner

CPGs are guidelines for best practice and are not intended as a substitute for good clinical judgment. Unusual patient presentations make it impossible to develop a CPG to match every possible clinical situation. The practitioner decides if a CPG should be applied based on patient assessment and the clinical impression. The practitioner must work in the best interest of the patient within the scope of practice for his/her clinical level on the PHECC Register. Consultation with fellow practitioners and or medical practitioners in challenging clinical situations is strongly advised.

The CPGs herein may be implemented provided:

- 1 The practitioner is in good standing on the PHECC Practitioner's Register.
- 2 The practitioner is acting on behalf of a licensed CPG provider (paid or voluntary).
- 3 The practitioner is privileged by the licensed CPG provider on whose behalf he/she is acting to implement the specific CPG.
- 4 The practitioner has received training on – and is competent in – the skills and medications specified in the CPG being utilised.

The medication dose specified on the relevant CPG shall be the definitive dose in relation to practitioner administration of medications. The principle of titrating the dose to the desired effect shall be applied. The onus rests on the practitioner to ensure that he/she is using the latest versions of CPGs which are available on the PHECC website www.phecc.ie

Definitions

Adult	A patient of 16 years or greater, unless specified on the CPG.
Child	A patient between 1 and less than or equal to (\leq) 15 years old, unless specified on the CPG
Infant	A patient between 4 weeks and less than 1 year old, unless specified on the CPG
Neonate	A patient less than 4 weeks old, unless specified on the CPG
Paediatric patient	Any child, infant or neonate

CPGs and the pre-hospital emergency care team

The aim of pre-hospital emergency care is to provide a comprehensive and coordinated approach to patient care management, thus providing each patient with the most appropriate care in the most efficient time frame.

In Ireland today, the provision of emergency care comes from a range of disciplines and includes responders (Cardiac First Responders, First Aid Responders and Emergency First Responders) and practitioners (Emergency Medical Technicians, Paramedics, Advanced Paramedics, Nurses and Doctors) from the statutory, private, auxiliary and voluntary services.

IMPLEMENTATION

CPGs set a consistent standard of clinical practice within the field of pre-hospital emergency care. By reinforcing the role of the practitioner, in the continuum of patient care, the chain of survival and the golden hour are supported in medical and traumatic emergencies respectively.

CPGs guide the practitioner in presenting to the acute hospital a patient who has been supported in the very early phase of injury/illness and in whom the danger of deterioration has lessened by early appropriate clinical care interventions.

CPGs presume no intervention has been applied, nor medication administered, prior to the arrival of the practitioner. In the event of another practitioner or responder initiating care during an acute episode, the practitioner must be cognisant of interventions applied and medication doses already administered and act accordingly.

In this care continuum, the duty of care is shared among all responders/practitioners of whom each is accountable for his/her own actions. The most qualified responder/practitioner on the scene shall take the role of clinical leader. Explicit handover between responders/practitioners is essential and will eliminate confusion regarding the responsibility for care.

In the absence of a more qualified practitioner, the practitioner providing care during transport shall be designated the clinical leader as soon as practical.

Emergency Medical Technician – Basic Tactical Emergency Care (EMT-BTEC)

EMT-BTEC certifies registered EMTs with additional knowledge and skill set for providing pre-hospital emergency care in hostile or austere environments. EMT-BTEC training is restricted to EMTs who have the potential to provide emergency care in hostile or austere environments and who are working or volunteering on behalf of a Licensed CPG Provider with specific approval for BTEC provision.

Emergency First Response – Basic Tactical Emergency Care (EFR-BTEC)

EFR-BTEC is a new education and training standard published in 2014. Persons certified at EFR-BTEC learn EFR and the additional knowledge and skill set for providing pre-hospital emergency care in hostile or austere environments. Entry to this course is restricted to people who have the potential to provide emergency first response in hostile or austere environments and who are working or volunteering on behalf of a Licensed CPG Provider with specific approval for BTEC provision.

First Aid Response

First Aid Response (FAR) is a new education and training standard published in 2014. This standard offers training and certification to individuals and groups who require a first aid skill set including cardiac first response. This standard is designed to meet basic first aid and basic life support (BLS) requirements that a certified person, known as a "First Aid Responder", may encounter in their normal daily activities.

Defibrillation Policy

The Medical Advisory Committee has recommended the following pre-hospital defibrillation policy;

- Advanced Paramedics should use manual defibrillation for all age groups.
- Paramedics may consider use of manual defibrillation for all age groups.
- EMTs and responders shall use AED mode for all age groups.

INDEX

EMERGENCY MEDICAL TECHNICIAN CPGs

SECTION 1 CARE PRINCIPLES	15
SECTION 2 PATIENT ASSESSMENT	16
Primary Survey Medical – Adult	16
Primary Survey Trauma – Adult	17
Secondary Survey Medical – Adult	18
Secondary Survey Trauma – Adult	19
Pain Management – Adult	20
SECTION 3 RESPIRATORY EMERGENCIES	21
Advanced Airway Management – Adult	21
Inadequate Ventilations – Adult	22
Exacerbation of COPD	23
Asthma - Adult	24
SECTION 4 MEDICAL EMERGENCIES	25
Basic Life Support – Adult	25
Foreign Body Airway Obstruction – Adult	26
VF or Pulseless VT – Adult	27
Asystole – Adult	28
Pulseless Electrical Activity – Adult	29
Post-Resuscitation Care – Adult	30
End of Life - DNR	31
Recognition of Death – Resuscitation not Indicated	32
Cardiac Chest Pain – Acute Coronary Syndrome	33
Symptomatic Bradycardia – Adult	34
Altered Level of Consciousness – Adult	35
Allergic Reaction/Anaphylaxis – Adult	36
Decompression Illness (DCI)	37
Epistaxis	38
Glycaemic Emergency – Adult	39
Hypothermia	40
Poisons – Adult	41
Seizure/Convulsion – Adult	42
Sepsis – Adult	43
Sickle Cell Crisis – Adult	44
Stroke	45
Mental Health Emergency	46
Behavioural Emergency	47
SECTION 5 OBSTETRIC EMERGENCIES	48
Pre-Hospital Emergency Childbirth	48
Basic Life Support – Neonate	49

INDEX

EMERGENCY MEDICAL TECHNICIAN CPGs

SECTION 6 TRAUMA	50
Burns – Adult	50
External Haemorrhage – Adult	51
Harness Induced Suspension Trauma	52
Head Injury – Adult	53
Heat-Related Emergency	54
Limb Injury – Adult	55
Shock from Blood Loss – Adult	56
Spinal Immobilisation – Adult	57
Submersion Incident	58
SECTION 7 PAEDIATRIC EMERGENCIES	59
Primary Survey Medical – Paediatric	59
Primary Survey Trauma – Paediatric	60
Secondary Survey – Paediatric	61
Pain Management – Paediatric	62
Inadequate Ventilations – Paediatric	63
Asthma - Paediatric	64
Stridor – Paediatric	65
Basic Life Support – Paediatric	66
Foreign Body Airway Obstruction – Paediatric	67
VF or Pulseless VT – Paediatric	68
Asystole/PEA – Paediatric	69
Symptomatic Bradycardia – Paediatric	70
Post Resuscitation Care – Paediatric	71
Allergic Reaction/Anaphylaxis – Paediatric	72
Glycaemic Emergency – Paediatric	73
Seizure/Convulsion – Paediatric	74
Pyrexia - Paediatric	75
Sickle Cell Crisis - Paediatric	76
External Haemorrhage – Paediatric	77
Shock from Blood Loss – Paediatric	78
Spinal Immobilisation – Paediatric	79
Burns – Paediatric	80
SECTION 8 PRE-HOSPITAL EMERGENCY CARE OPERATIONS	81
Major Emergency – First Practitioners on site	81
Major Emergency – Operational Control	82
Triage Sieve	83

CLINICAL PRACTICE GUIDELINES for EMERGENCY MEDICAL TECHNICIAN

(CODES EXPLANATION)



Emergency Medical Technician
(Level 4) for which the CPG pertains



Paramedic
(Level 5) for which the CPG pertains



Advanced Paramedic
(Level 6) for which the CPG pertains




An **EMT** who has completed Basic Tactical Emergency Care training and has been privileged to operate in adverse conditions




A parallel process
Which may be carried out in parallel with other sequence steps




A cyclical process in which a number of sequence steps are completed




A sequence (skill) to be performed




A mandatory sequence (skill) to be performed



A decision process
The Practitioner must follow one route



Given the clinical presentation consider the treatment option specified



Finding following clinical assessment, leading to treatment modalities



Reassess the patient following intervention



Contact Ambulance Control and request Advanced Life Support (AP or doctor)



Consider requesting an ALS response, based on the clinical findings




Emergency Medical Technician or lower clinical levels not permitted this route



Transport to an appropriate medical facility and maintain treatment en-route




Transport to an appropriate medical facility and maintain treatment en-route, if having contacted Ambulance Control there is no ALS available



An instruction box for information




Special instructions
Which the Practitioner must follow



A skill or sequence that only pertains to Paramedic or higher clinical levels




Special authorisation
This authorises the Practitioner to perform an intervention under specified conditions




CPG numbering system


4/5/6 = clinical levels to which the CPG pertains
x = section in CPG manual, y = CPG number in sequence
mm/yy = month/year CPG published




Consider requesting a Paramedic response, based on the clinical findings



A medication which may be administered by an EMT or higher clinical level
The medication name, dose and route is specified



A medication which may be administered by a Paramedic or higher clinical level
The medication name, dose and route is specified



A medication which may be administered by an Advanced Paramedic
The medication name, dose and route is specified



A direction to go to a specific CPG following a decision process
Note: only go to the CPGs that pertain to your clinical level



A clinical condition that may precipitate entry into the specific CPG

SECTION 1

CARE PRINCIPLES

Care principles are goals of care that apply to all patients. Scene safety, standard precautions, patient assessment, primary and secondary surveys and the recording of interventions and medications on the Patient Care Report (PCR) or the Ambulatory Care Report (ACR) are consistent principles throughout the guidelines and reflect the practice of practitioners. Care principles are the foundations for risk management and the avoidance of error.


PHECC Care Principles

- 1 Ensure the safety of yourself, other emergency service personnel, your patients and the public.
- 2 Seek consent prior to initiating interventions and/or administering medications.
- 3 Identify and manage life-threatening conditions.
- 4 Ensure adequate ventilation and oxygenation.
- 5 Optimise tissue perfusion.
- 6 Provide appropriate pain relief.
- 7 Identify and manage other conditions.
- 8 Place the patient in the appropriate posture according to the presenting condition.
- 9 Ensure the maintenance of normal body temperature (unless a CPG indicates otherwise).
- 10 Provide reassurance at all times.
- 11 Monitor and record patient's vital observations.
- 12 Maintain responsibility for patient care until handover to an appropriate practitioner.
- 13 Arrange transport to an appropriate medical facility as necessary and in an appropriate time frame.
- 14 Complete patient care records following an interaction with a patient.
- 15 Identify the clinical leader on scene; this shall be the most qualified practitioner on scene. In the absence of a more qualified practitioner, the practitioner providing care during transport shall be designated the clinical leader as soon as practical.

SECTION 2 PATIENT ASSESSMENT

4/5/6.2.1
Version 3, 02/14


Primary Survey Medical – Adult




SECTION 2 PATIENT ASSESSMENT

4/5/6.2.2
Version 3, 02/14

Primary Survey Trauma – Adult



The primary survey is focused on establishing the patient's clinical status and only applying interventions when they are essential to maintain life. It should be completed within one minute of arrival on scene.




Reference: ILCOR Guidelines 2010

SECTION 2
PATIENT ASSESSMENT

4.2.4
Version 2, 09/11

Secondary Survey Medical – Adult

EMT




Reference: Sanders, M. 2001, Paramedic Textbook 2nd Edition, Mosby
Gleadle, J. 2003, History and Examination at a glance, Blackwell Science
Rees, JE, 2003, Early Warning Scores, World Anaesthesia Issue 17, Article 10

SECTION 2
PATIENT ASSESSMENT

4.2.5
Version 1, 05/08

Secondary Survey Trauma – Adult

EMT



<p>Markers for multi-system trauma Systolic BP < 90 Respiratory rate < 10 or > 29 Heart rate > 120 AVPU = V, P or U on scale Mechanism of Injury</p>
--


Reference: McSwain, N. et al, 2003, PHTLS Basic and advanced prehospital trauma life support, 5th Edition, Mosby

SECTION 2
PATIENT ASSESSMENT


4/5/6.2.6
Version 4, 02/14

Pain Management – Adult

EMT P
AP



Analogue Pain Scale
0 = no pain.....10 = unbearable



Special Authorisation:
AP APs are authorised to administer Morphine, up to 10 mg IM, if IV not accessible, the patient is cardiovascularly stable and no cardiac chest pain present


Reference: World Health Organization, Pain Ladder

SECTION 3
RESPIRATORY EMERGENCIES

4.3.1
Version 3, 07/13

Advanced Airway Management – Adult

CFR - A EMT



Minimum interruptions of chest compressions.
Maximum hands off time 10 seconds.


Maintain adequate ventilation and oxygenation throughout procedures

Following successful Advanced Airway management:-
i) Ventilate at 8 to 10 per minute.
ii) Unsynchronised chest compressions continuous at 100 to 120 per minute


Equipment list
Non-inflatable supraglottic airway

EMT Special Authorisation:
EMTs may use cuffed supraglottic airways subject to maintaining competence and Medical Director authorisation

SECTION 3
RESPIRATORY EMERGENCIES



SECTION 3
RESPIRATORY EMERGENCIES



An exacerbation of COPD is defined as:
An event in the natural course of the disease characterised by a change in the patient's baseline dyspnoea, cough and/or sputum beyond day-to-day variability sufficient to warrant a change in management. (European Respiratory Society)

SECTION 3
RESPIRATORY EMERGENCIES




Reference: HSE National Asthma Programme 2012, Emergency Asthma Guidelines, British Thoracic Society, 2008, British Guidelines on the Management of Asthma, a national clinical guideline

SECTION 4
MEDICAL EMERGENCIES

4/5/6.4.1
Version 2, 06/11

Basic Life Support – Adult



Initiate mobilisation of 3 to 4 practitioners / responders on site to assist with cardiac arrest management

Cardiac Arrest

Request
ALS


Attach defibrillation pads
Commence CPR while defibrillator is being prepared only if 2nd person available
30 Compressions : 2 ventilations.
Oxygen therapy

Chest compressions
Rate: 100 to 120/ min
Depth: at least 5 cm

Ventilations
Rate: 10/ min (1 every 6 sec)
Volume: 500 to 600 mL

AP Change defibrillator to manual mode
P Consider changing defibrillator to manual mode


Continue CPR while defibrillator is charging



Immediately resume CPR x 2 minutes

Minimum interruptions of chest compressions.
Maximum hands off time 10 seconds.

Rhythm check *



If an Implantable Cardioverter Defibrillator (ICD) is fitted in the patient treat as per CPG. It is safe to touch a patient with an ICD fitted even if it is firing.

* +/- Pulse check: pulse check after 2 minutes of CPR if potentially perfusing rhythm


Reference: ILCOR Guidelines 2010

SECTION 4
MEDICAL EMERGENCIES

4/5.4.2
Version 1, 05/08


Foreign Body Airway Obstruction – Adult

EMT P




After each cycle of CPR open mouth and look for object. If visible attempt once to remove it

SECTION 4
MEDICAL EMERGENCIES




SECTION 4
MEDICAL EMERGENCIES



* +/- Pulse check: pulse check after 2 minutes of CPR if potentially perfusing rhythm

Reference: ILCOR Guidelines 2010

SECTION 4
MEDICAL EMERGENCIES



* +/- Pulse check: pulse check after 2 minutes of CPR if potentially perfusing rhythm

Reference: ILCOR Guidelines 2010


SECTION 4
MEDICAL EMERGENCIES

4.4.7
Version 2, 03/11

Post-Resuscitation Care – Adult


EMT

Equipment list
Cold packs



Titrate O₂ to 94% - 98%

For active cooling place cold packs at arm pit, groin & abdomen



Reference: ILCOR Guidelines 2010


SECTION 4

MEDICAL EMERGENCIES

4.4.8
Version 1, 06/10

End of Life – DNR

EMT




SECTION 4
MEDICAL EMERGENCIES

4.4.9
Version 1, 05/08

Recognition of Death – Resuscitation not Indicated

EMT




SECTION 4
MEDICAL EMERGENCIES

4.4.10
Version 2, 09/11

Cardiac Chest Pain – Acute Coronary Syndrome

EMT



SECTION 4
MEDICAL EMERGENCIES


4/5/6.4.11
Version 2, 02/14

Symptomatic Bradycardia – Adult

EMT P
AP

Symptomatic includes;
Acute altered mental status
Ischemic chest discomfort
Acute heart failure
Hypotension
Signs of shock

Titrate Atropine to
effect (HR > 60)




SECTION 4
MEDICAL EMERGENCIES

4.4.14
Version 1, 05/08

Altered Level of Consciousness – Adult

EMT




SECTION 4
MEDICAL EMERGENCIES

4.4.15
Version 3, 02/14

Allergic Reaction/Anaphylaxis – Adult

EMT



Nebulised Salbutamol may be substituted with up to 5 puffs of Salbutamol aerosol

Mild
Urticaria and/or angio oedema

Moderate
Mild symptoms + simple bronchospasm


Severe/ anaphylaxis
Moderate symptoms + haemodynamic and or respiratory compromise

SECTION 4
MEDICAL EMERGENCIES

4/5/6.4.16
Version 2, 07/11

Decompression Illness (DCI)

EMT P
AP



Special Authorisation:
Paramedics are authorised to continue the established infusion in the absence of an Advanced Paramedic or Doctor during transportation


Reference: The Primary Clinical Care Manual 3rd Edition, 2003, Queensland Health and the Royal Flying Doctor Service (Queensland Section)

SECTION 4
MEDICAL EMERGENCIES

4/5/6.4.17
Version 2, 01/13

Epistaxis

EMT P
AP




Reference: Management of Acute Epistaxis 2011, Ola Bamimore, MD; Chief Editor: Steven C Dronen, MD, <http://emedicine.medscape.com/article/764719-overview#showall>

SECTION 4
MEDICAL EMERGENCIES

4.4.19
Version 1, 05/08

Glycaemic Emergency – Adult

EMT




SECTION 4
MEDICAL EMERGENCIES

4.4.21
Version 2, 02/14

Hypothermia

EMT




Reference: Golden, F & Tipton M, 2002, Essentials of Sea Survival, Human Kinetics
 AHA, 2005, Part 10.4: Hypothermia, Circulation 2005;112:136-138
 Soar, J et al, 2005, European Resuscitation Council Guidelines for Resuscitation 2005, Section 7. Cardiac arrest in special circumstances, Resuscitation (2005) 6751, S135-S170
 Pennington M, et al, 1994, Wilderness EMT, Wilderness EMS Institute

SECTION 4
MEDICAL EMERGENCIES

4/5.4.22
Version 3, 02/14

Poisons – Adult

EMT P



Reference:

ILCOR Guidelines 2010

Boyer, E. 2012, Management of Opioid Analgesic Overdose, N Engl J Med 2012;367:146-55.DOI: 10.1056/NEJMra1202561


SECTION 4
MEDICAL EMERGENCIES

4.4.23
Version 2, 07/11

Seizure/Convulsion – Adult

EMT



Consider other causes of seizures
Meningitis
Head injury
Hypoglycaemia
Eclampsia
Fever
Poisons
Alcohol/drug withdrawal



SECTION 4 MEDICAL EMERGENCIES


4/5/6.4.24
Version 3, 02/14

Sepsis – Adult



Special Authorisation:
Paramedics are authorised to continue the established infusion in the absence of an Advanced Paramedic or Doctor during transportation

SECTION 4
MEDICAL EMERGENCIES



Special Authorisation:
Paramedics are authorised to continue the established infusion in the absence of an Advanced Paramedic or Doctor during transportation


Reference: Rees, D, 2003, GUIDELINES FOR THE MANAGEMENT OF THE ACUTE PAINFUL CRISIS IN SICKLE CELL DISEASE; British Journal of Haematology, 2003, 120, 744-752

SECTION 4
MEDICAL EMERGENCIES

4.4.28
Version 1, 05/08

Stroke

EMT



Oxygen therapy
Maintain SpO₂ between 94% to 98%
(lower range if COPD)

F – facial weakness
Can the patient smile?, Has their mouth or eye drooped? Which side?
A – arm weakness
Can the patient raise both arms and maintain for 5 seconds?
S – speech problems
Can the patient speak clearly and understand what you say?
T – time to transport now if FAST positive

SECTION 4
MEDICAL EMERGENCIES

4/5.4.29
Version 1, 05/08

Mental Health Emergency

EMT P

RMP – Registered Medical Practitioner
RPN – Registered Psychiatric Nurse

Practitioners may not compel a patient to accompany them or prevent a patient from leaving an ambulance vehicle

If potential to harm self or others ensure minimum two people accompany patient in saloon of ambulance at all times

Behaviour abnormal with previous psychiatric history

RMP or RPN in attendance or have made arrangements for voluntary/assisted admission

Co-operate as appropriate with medical or nursing team

Transport patient to an Approved Centre

Potential to harm self or others

Request control to inform Gardai

Reassure patient
Explain what is happening at all times
Avoid confrontation

Attempt verbal de-escalation

Combative with hallucinations or Paranoia & risk to self or others

Request ALS

Patient agrees to travel

Request as appropriate
- Gardai
- Medical Practitioner
- Mental health team

Aid to Capacity Evaluation
1. Patient verbalises/ communicates understanding of clinical situation?
2. Patient verbalises/ communicates appreciation of applicable risk?
3. Patient verbalises/ communicates ability to make alternative plan of care?
If no to any of the above consider Patient Incapacity



Reference: Reference Guide to the Mental Health Act 2001, Mental Health Commission
HSE Mental Health Services

SECTION 4

MEDICAL EMERGENCIES

4/5/6.4.30
Version 1, 05/08


Behavioural Emergency

EMT P

AP

Practitioners may not compel a patient to accompany them or prevent a patient from leaving an ambulance vehicle

If potential to harm self or others ensure minimum two people accompany patient in saloon of ambulance at all times



Aid to Capacity Evaluation

1. Patient verbalises/ communicates understanding of clinical situation?
2. Patient verbalises/ communicates appreciation of applicable risk?
3. Patient verbalises/ communicates ability to make alternative plan of care?

If no to any of the above consider Patient Incapacity


Reference: HSE Mental Health Services

SECTION 5
OBSTETRIC EMERGENCIES

4.5.1
Version 1, 05/08

Pre-Hospital Emergency Childbirth

EMT




SECTION 5
OBSTETRIC EMERGENCIES

4.5.2
Version 1, 05/08

Basic Life Support – Neonate (< 4 weeks)



EMT



SECTION 6 TRAUMA


4/5/6.6.1
Version 2, 07/11

Burns – Adult




Reference: Allison, K et al, 2004, Consensus on the prehospital approach to burns patient management, Emerg Med J 2004; 21:112-114
Sanders, M, 2001, Paramedic Textbook 2nd Edition, Mosby

SECTION 6 TRAUMA



SECTION 6
TRAUMA




Reference:
Adish A et al, 2009, Evidence-based review of the current guidance on first aid measures for suspension trauma, Health and Safety Executive (UK) Research report RR708
Australian Resuscitation Council, 2009, Guideline 9.1.5 Harness Suspension Trauma first aid management.
Thomassen O et al, Does the horizontal position increase risk of rescue death following suspension trauma?, *Emerg Med J* 2009;26:896-898 doi:10.1136/emj.2008.064931

SECTION 6
TRAUMA

4.6.5
Version 2, 01/13

Head Injury – Adult

EMT



Equipment list
Extrication device
Long board
Vacuum mattress
Orthopaedic stretcher
Rigid cervical collar


Reference;
Mc Swain, N, 2011, PHTLS Prehospital Trauma Life Support 7th Edition, Mosby

SECTION 6 TRAUMA

4/5/6.6.6
Version 1, 12/13

Heat-Related Emergency – Adult

EMT P
AP




Special Authorisation:
Paramedics are authorised to continue the established infusion in the absence of an Advanced Paramedic or Doctor during transportation

SECTION 6 TRAUMA

4/5/6.6.7
Version 4, 02/14

Limb Injury – Adult

EMT P
AP



Equipment list
Traction splint
Box splint
Frac straps
Triangular bandages
Vacuum splints
Long board
Orthopaedic stretcher
Cold packs
Elastic bandages
Pelvic splinting device

Contraindications for application of traction splint
1 # pelvis
2 # knee
3 Partial amputation
4 Injuries to lower third of lower leg
5 Hip injury that prohibits normal alignment

For a limb-threatening injury treat as an emergency and pre alert ED


Reference: An algorithm guiding the evaluation and treatment of acute primary patellar dislocations, Mehta VM et al. Sports Med Arthrosc. 2007 Jun;15(2):78-81

SECTION 6
TRAUMA

4.6.8
Version 2, 01/13

Shock from Blood Loss – Adult

EMT



<p>Signs of poor perfusion A: (Not affected) B: Tachypnea C: Tachycardia Delayed capillary refill Diminished/absent peripheral pulses D: V, P or U / Irritability / confusion E: Cool, pale & moist skin</p>
--

SECTION 6
TRAUMA

4.6.9
Version 1, 05/08


Spinal Immobilisation – Adult

EMT

If in doubt,
treat as
spinal injury


Do not forcibly restrain a
patient that is combative

Equipment list
Long board
Vacuum mattress
Orthopaedic stretcher
Rigid cervical collar



Dangerous mechanism include;
 Fall ≥ 1 metre/ 5 steps
 Axial load to head
 MVC > 100 km/hr, rollover or ejection
 ATV collision
 Bicycle collision
 Pedestrian v vehicle

SECTION 6 TRAUMA




Reference: Golden, F & Tipton M, 2002, Essentials of Sea Survival, Human Kinetics
 Verie, M, 2007, Near Drowning, E medicine, www.emedicine.com/ped/topic20570.htm
 Shepherd, S, 2005, Submersion Injury, Near Drowning, E Medicine, www.emedicine.com/emerg/topic744.htm
 AHA, 2005, Part 10.3: Drowning, Circulation 2005;112:133-135
 Soar, J et al, 2005, European Resuscitation Council Guidelines for Resuscitation 2005, Section 7. Cardiac arrest in special circumstances, Resuscitation (2005) 6751, S135-S170

SECTION 7 PAEDIATRIC EMERGENCIES


4/5/6.7.1
Version 4, 12/13

Primary Survey Medical – Paediatric (≤ 15 Years)

EMT P AP



The primary survey is focused on establishing the patient's clinical status and only applying interventions when they are essential to maintain life. It should be completed within one minute of arrival on scene.



Ref: Pediatric Education for Prehospital Professionals

Consider Oxygen therapy

Go to Secondary Survey CPG

If child protection concerns are present

Report findings as per Children First guidelines to ED staff and line manager in a confidential manner


Normal ranges		
Age	Pulse	Respirations
Infant	100 – 160	30 – 60
Toddler	90 – 150	24 – 40
Pre school	80 – 140	22 – 34
School age	70 – 120	18 – 30

Reference:
ILCOR Guidelines 2010, American Academy of Pediatrics, 2000, Pediatric Education for Prehospital Professionals
Department of Children and Youth Affairs, 2011, Children First: National Guidance for the Protection and Welfare of Children



SECTION 7 PAEDIATRIC EMERGENCIES

4/5/6.7.2
Version 4, 12/13

Primary Survey Trauma – Paediatric (≤ 15 years)



The primary survey is focused on establishing the patient's clinical status and only applying interventions when they are essential to maintain life. It should be completed within one minute of arrival on scene.





Normal ranges		
Age	Pulse	Respirations
Infant	100 – 160	30 – 60
Toddler	90 – 150	24 – 40
Pre school	80 – 140	22 – 34
School age	70 – 120	18 – 30

Reference: ILCOR Guidelines 2010, American Academy of Pediatrics, 2000, Pediatric Education for Prehospital Professionals, Department of Children and Youth Affairs, 2011, Children First, National Guidance for the Protection and Welfare of Children

SECTION 7
PAEDIATRIC EMERGENCIES

4/5/6.7.4
Version 3, 12/13

Secondary Survey – Paediatric (≤ 15 years)



Use age appropriate language for patient



Children and adolescents should always be examined with a chaperone (usually a parent) where possible

Estimated weight
Neonate = 3.5 Kg
Six months = 6 Kg
One to five years = (age x 2) + 8 Kg
Greater than 5 years = (age x 3) + 7 Kg

Normal ranges		
Age	Pulse	Respirations
Infant	100 – 160	30 – 60
Toddler	90 – 150	24 – 40
Pre school	80 – 140	22 – 34
School age	70 – 120	18 – 30


Reference:
Miall, Lawrence et al, 2003, Paediatrics at a Glance, Blackwell Publishing
Department of Children and Youth Affairs, 2011, Children First: National Guidance for the Protection and Welfare of Children
Luscombe, M et al 2010, BMJ, Weight estimation in paediatrics: a comparison of the APLS formula and the formula 'Weight=(age)+7'

SECTION 7
PAEDIATRIC EMERGENCIES




Decisions to give analgesia must be based on clinical assessment and not directly on a linear scale

SECTION 7
PAEDIATRIC EMERGENCIES



SECTION 7
PAEDIATRIC EMERGENCIES





Reference: HSE National Asthma Programme 2012, Emergency Asthma Guidelines, British Thoracic Society, 2008, British Guidelines on the Management of Asthma, a national clinical guideline

SECTION 7
PAEDIATRIC EMERGENCIES

4/5/6.7.13
Version 2, 12/13


Stridor – Paediatric (≤ 15 years)



SECTION 7
PAEDIATRIC EMERGENCIES

4/5/6.7.20
Version 2, 12/13

Basic Life Support – Paediatric (≤ 15 Years)



Initiate mobilisation of 3 to 4 practitioners / responders on site to assist with cardiac arrest management

Cardiac arrest
or
pulse < 60 per minute with signs of poor perfusion

Give 5 rescue ventilations
Oxygen therapy

Request
ALS

< 8 years use paediatric defibrillation system (if not available use adult pads)

Minimum interruptions of chest compressions.
Maximum hands off time 10 seconds.

One rescuer CPR 30 : 2
Two rescuer CPR 15 : 2
Compressions : Ventilations

Commence chest Compressions
Continue CPR (30:2) until defibrillator is attached

Chest compressions
Rate: 100 to 120/ min
Depth: 1/3 depth of chest
Child; two hands
Small child; one hand
Infant (< 1); two fingers

Yes < 8 years No

AP Change defibrillator to manual mode
P Consider changing defibrillator to manual mode

Apply paediatric system AED pads

Apply adult defibrillation pads

With two rescuer CPR use two thumb-encircling hand chest compression for infants

Shockable VF or pulseless VT Non - Shockable Asystole or PEA

Continue CPR while defibrillator is charging

Give 1 shock

Immediately resume CPR x 2 minutes

Rhythm check *

Go to VF / Pulseless VT CPG

Go to Post Resuscitation Care CPG

Asystole / PEA

Go to Asystole / PEA CPG

Infant AED
It is extremely unlikely to ever have to defibrillate a child less than 1 year old. Nevertheless, if this were to occur the approach would be the same as for a child over the age of 1. The only likely difference being, the need to place the defibrillation pads anterior (front) and posterior (back), because of the infant's small size.

* +/- Pulse check: pulse check after 2 minutes of CPR if potentially perfusing rhythm


Reference: ILCOR Guidelines 2010

SECTION 7
PAEDIATRIC EMERGENCIES

4/5.7.21
Version 2, 12/13


Foreign Body Airway Obstruction – Paediatric (≤ 15 years)

EMT P




After each cycle of CPR open mouth and look for object. If visible attempt once to remove it

SECTION 7
PAEDIATRIC EMERGENCIES



SECTION 7
PAEDIATRIC EMERGENCIES



* +/- Pulse check: pulse check after 2 minutes of CPR if potentially perfusing rhythm


Reference: ILCOR Guidelines 2010

SECTION 7
PAEDIATRIC EMERGENCIES

4/5/6.7.24
Version 3, 10/13

Symptomatic Bradycardia – Paediatric (≤ 15 years)

EMT P
AP



Reference: International Liaison Committee on Resuscitation, 2010, Part 6: Paediatric basic and advanced life support, Resuscitation (2005) 67, 271 – 291

SECTION 7
PAEDIATRIC EMERGENCIES


4.7.25
Version 2, 12/13

Post-Resuscitation Care – Paediatric (≤ 15 years)

EMT

- Equipment list
- Cold packs

Titrate O₂ to 96% - 98%



For active cooling place cold packs at arm pit, groin & abdomen




Reference: ILCOR Guidelines 2010

SECTION 7
PAEDIATRIC EMERGENCIES

4.7.31
Version 4, 02/14

Allergic Reaction/Anaphylaxis – Paediatric (≤ 15 years)

EMT



Salbutamol NEB may be substituted with Salbutamol aerosol 0.1 mg. If no improvement Salbutamol may be repeated; for < 5 year olds up to 3 times, for ≥ 5 year olds up to 5 times, prn

Mild
Urticaria and/or angio oedema

Moderate
Mild symptoms + simple bronchospasm


Severe
Moderate symptoms + haemodynamic and/or respiratory compromise

SECTION 7
PAEDIATRIC EMERGENCIES

4.7.32
Version 3, 12/13

Glycaemic Emergency – Paediatric (≤ 15 years)

EMT




SECTION 7
PAEDIATRIC EMERGENCIES

4.7.33
Version 3, 02/14

Seizure/Convulsion – Paediatric (≤ 15 years)

EMT



Consider other causes of seizures
Meningitis
Head injury
Hypoglycaemia
Fever
Poisons
Alcohol/drug withdrawal



SECTION 7
PAEDIATRIC EMERGENCIES

4/5/6.7.35
Version 1, 12/13

Pyrexia – Paediatric (≤ 15 years)




Reference: ILCOR Guidelines 2010
RFDS, 2011, Primary Clinical Care Manual

SECTION 7
PAEDIATRIC EMERGENCIES

4/5/6.7.36
Version 1, 12/13

Sickle Cell Crisis – Paediatric (≤ 15 years)

EMT P
AP




Consider patient's care plan

P Special Authorisation:
Paramedics are authorised to continue the established infusion in the absence of an Advanced Paramedic or Doctor during transportation

Reference: Rees, D, 2003, GUIDELINES FOR THE MANAGEMENT OF THE ACUTE PAINFUL CRISIS IN SICKLE CELL DISEASE; British Journal of Haematology, 2003, 120, 744–752

SECTION 7
PAEDIATRIC EMERGENCIES




Reference:
ILCOR Guidelines 2010,
Granville-Chapman J, et al. Pre-hospital haemostatic dressings: A systematic review. Injury (2010), doi: 10.1016/j.injury.2010.09.037

SECTION 7
PAEDIATRIC EMERGENCIES

4.7.51
Version 2, 12/13


Shock from Blood Loss – Paediatric (≤ 15 years)

EMT



Signs of inadequate perfusion
A: (not directly affected)
B: Increased respiratory rate (without increased effort)
C: Tachycardia
Diminished/absent peripheral pulses
Delayed capillary refill
D: Irritability/ confusion / ALoC
E: Cool extremities, mottling

SECTION 7 PAEDIATRIC EMERGENCIES




Paediatric spinal injury indications include

- Pedestrian v auto
- Passenger in high speed vehicle collision
- Ejection from vehicle
- Sports/ playground injuries
- Falls from a height
- Axial load to head

Do not forcibly restrain a patient that is combative

References:
Viccello, P, et al, 2001, A Prospective Multicentre Study of Cervical Spine Injury in Children, Pediatrics vol 108, e20
Slack, S. & Clancy, M, 2004, Clearing the cervical spine of paediatric trauma patients, EMJ 21; 189-193

SECTION 7
PAEDIATRIC EMERGENCIES




Reference: Allison, K et al, 2004, Consensus on the prehospital approach to burns patient management, Emerg Med J 2004; 21:112-114
Sanders, M, 2001, Paramedic Textbook 2nd Edition, Mosby


SECTION 8
PRE-HOSPITAL EMERGENCY CARE OPERATIONS

4/5/6.8.1
Version 2, 01/13

Major Emergency (Major Incident) – First Practitioners on site



Irish (Major Emergency) terminology in black
UK (Major Incident) terminology in blue



If single Practitioner is first on site combine both roles until additional Practitioners arrive

METHANE message
 M – Major Emergency declaration / standby
 E – Exact location of the emergency
 T – Type of incident (transport, chemical etc.)
 H – Hazards present and potential
 A – Access / egress routes
 N – Number of casualties (injured or dead)
 E – Emergency services present and required

The first ambulance crew does not provide care or transport of patients as this interferes with their ability to liaise with other services, to assess the scene and to provide continuous information as the incident develops


The principles and terminology of Major Incident Medical Management and Support (MIMMS) has been used with the kind permission of the Advanced Life Support Group, UK

Reference: A Framework for Major Emergency Management, 2006, Inter-Departmental Committee on Major Emergencies (Replaced by National Steering Group on Major Emergency Management)

SECTION 8
PRE-HOSPITAL EMERGENCY CARE OPERATIONS


4/5/6.8.2
Version 2, 01/13

Major Emergency (Major Incident) – Operational Control



Irish (Major Emergency) terminology in black
UK (Major Incident) terminology in blue

If Danger Area identified, entry to Danger Area is controlled by a Senior Fire Officer or an Garda Síochána



Entry to Inner Cordon (Bronze Area) is limited to personnel providing emergency care and or rescue
Personal Protective Equipment required

Entry to Outer Cordon (Silver area) is controlled by an Garda Síochána

Management structure for;
Outer Cordon, Tactical Area (Silver Area)
On-Site Co-ordinator
HSE Controller of Operations (Ambulance Incident Officer)
Site Medical Officer (Medical Incident Officer)
Local Authority Controller of Operations (Fire Incident Officer)
Garda Controller of Operations (Police Incident Officer)

Management structure for;
Inner Cordon, Operational Area (Bronze Area)
Forward Ambulance Incident Officer (Forward Ambulance Incident Officer)
Forward Medical Incident Officer (Forward Medical Incident Officer)
Fire Service Incident Commander (Forward Fire Incident Officer)
Garda Cordon Control Officer (Forward Police Incident Officer)

Please note that Controller of Operations may be other than ambulance or fire officers, depending on the nature of the emergency

Other management functions for;
Major Emergency site
Casualty Clearing Officer
Triage Officer
Ambulance Parking Point Officer
Ambulance Loading Point Officer
Communications Officer
Safety Officer





Reference: A Framework for Major Emergency Management, 2006, Inter-Departmental Committee on Major Emergencies (Replaced by National Steering Group on Major Emergency Management)
The principles and terminology of Major Incident Medical Management and Support (MIMMS) has been used with the kind permission of the Advanced Life Support Group, UK

SECTION 8
PRE-HOSPITAL EMERGENCY CARE OPERATIONS

4/5/6.8.3
Version 1, 05/08

Triage Sieve



Triage is a dynamic process

The principles and terminology of Major Incident Medical management and Support (MIMMS) has been used with the kind permission of the Advanced Life Support Group, UK

APPENDIX 1 MEDICATION FORMULARY

The Medication Formulary is published by the Pre-Hospital Emergency Care Council (PHECC) to enable pre-hospital emergency care practitioners to be competent in the use of medications permitted under the Medicinal Products 7th Schedule (SI 300 of 2014). This is a summary document only and practitioners are advised to consult with official publications to obtain detailed information about the medications used.

The Medication Formulary is recommended by the Medical Advisory Committee (MAC) prior to publication by Council.

The medications herein may be administered provided:

- 1 The practitioner is in good standing on the PHECC practitioner's Register.
- 2 The practitioner complies with the Clinical Practice Guidelines (CPGs) published by PHECC.
- 3 The practitioner is acting on behalf of an organisation (paid or voluntary) that is a PHECC licensed CPG provider.
- 4 The practitioner is privileged, by the organisation on whose behalf he/she is acting, to administer the medications.
- 5 The practitioner has received training on, and is competent in, the administration of the medication.
- 6 The medications are listed on the Medicinal Products 7th Schedule.

The context for administration of the medications listed here is outlined in the CPGs.

Every effort has been made to ensure accuracy of the medication doses herein. The dose specified on the relevant CPG shall be the definitive dose in relation to practitioner administration of medications. The principle of titrating the dose to the desired effect shall be applied. The onus rests on the practitioner to ensure that he/she is using the latest versions of CPGs which are available on the PHECC website www.phecc.ie

Sodium Chloride 0.9% (NaCl) is the IV/IO fluid of choice for pre-hospital emergency care.

Water for injection shall be used when diluting medications, however if not available NaCl (0.9%) may be used if not contraindicated.

All medication doses for patients' ≤ 15 years shall be calculated on a weight basis unless an age-related dose is specified for that medication.

The route of administration should be appropriate to the patients clinical presentation. IO access is authorised for Advanced Paramedics for life threatening emergencies (or under medical direction).

The dose for paediatric patients may never exceed the adult dose.

Paediatric weight estimations acceptable to PHECC are:

Neonate	3.5 Kg
Six months	6 Kg
One to five years	(age x 2) + 8 Kg
Greater than 5 years	(age x 3) + 7 Kg

Reviewed on behalf of PHECC by Prof Peter Weedle, Adjunct Professor of Clinical Pharmacy, School of Pharmacy, University College Cork.

This version contains 11 medications.

APPENDIX 1 MEDICATION FORMULARY

Amendments to the 2012 Edition

The paediatric age range has been increased to reflect the HSE National Clinical Programme for Paediatrics and Neonatology age profile:

A paediatric patient is defined as a patient up to the eve of his/her 16th birthday (≤ 15 years).


Water for injection shall be used when diluting medications, however if not available NaCl (0.9%) may be used if not contraindicated.

The paediatric weight estimation formulae have been modified.


New Medications introduced;

- Ibuprofen
- Naloxone

Epinephrine (1:1,000)		
HEADING	ADD	DELETE
Usual Dosages	Auto-injector	EpiPen® Jr

Ibuprofen		
HEADING	ADD	DELETE
Clinical Level		
Presentation	400 mg tablet	
Description	It is an anti-inflammatory analgesic	It is used to reduce mild to moderate pain
Additional information	Caution with significant burns or poor perfusion due to risk of kidney failure Caution if concurrent NSAIDs use	

APPENDIX 1
MEDICATION FORMULARY

Naloxone		
HEADING	ADD	DELETE
Clinical level		
Administration	Intranasal (IN). CPG: 6.4.23, 4/5.4.23, 4/5/6.7.5	CPG: 5/6.3.2, 5/6.7.5
Indications	Inadequate respiration and/or ALoC following known or suspected narcotic overdose	Respiratory rate < 10 secondary to known or suspected narcotic overdose
Usual Dosages	Adult: 0.8 mg (800 mcg) IN (EMT) Paediatric: 0.02 mg/Kg (20 mcg/Kg) IN (EMT)	(Paramedic repeats by one prn)

Nitrous Oxide 50% and Oxygen 50% (Entonox®)		
HEADING	ADD	DELETE
Additional information	Caution when using Entonox for greater than one hour for Sickle Cell Crisis	

Oxygen		
HEADING	ADD	DELETE
Contraindications		Paraquat poisoning
Indications	Sickle Cell Disease - 100%	
Additional Information	Caution with paraquat poisoning, administer oxygen if SpO ₂ < 92%	

APPENDIX 1 MEDICATION FORMULARY

Paracetamol		
HEADING	ADD	DELETE
Presentation	250 mg in 5 mL	
Indications	Pyrexia	Pyrexia following seizure for paediatric patients. Advanced Paramedics may administer Paracetamol, in the absence of a seizure for the current episode, provided the paediatric patient is pyrexial and has a previous history of febrile convulsions.
Contraindications	< 1 month old	
Usual Dosages	> 1 month < 1 year - 90 mg PR.	< 1 year - 60 mg PR

Salbutamol		
HEADING	ADD	DELETE
Administration		Advanced Paramedics may repeat Salbutamol x 3
Usual Dosages	<p>Adult: .. (or 0.1 mg metered aerosol spray x 5) Repeat at 5 min prn (EFRs: 0.1 mg metered aerosol spray x 2)</p> <p>Paediatric: < 5 yrs...(or 0.1 mg metered aerosol spray x 3) ≥ 5 yrs...(or 0.1 mg metered aerosol spray x 5) Repeat at 5 min prn (EFRs: 0.1 mg metered aerosol spray x 2)</p>	<p>Adult: Repeat at 5 min prn (APs x 3 and Ps x 1) (EMTs & EFRs: 0.1 mg metered aerosol spray x 2)</p> <p>Paediatric: Repeat at 5 min prn (APs x 3 and Ps x 1) (EMTs & EFRs: 0.1 mg metered aerosol spray x 2)</p>

Please visit www.phecc.ie for the latest edition/version.

APPENDIX 1 MEDICATION FORMULARY

LIST OF MEDICATIONS

Aspirin	89
Epinephrine 1mg/1ml (1:1000)	90
Glucagon	91
Glucose gel	92
Glyceryl Trinitrate (GTN)	93
Ibuprofen	94
Naloxone	95
Nitrous Oxide 50% and Oxygen 50% (Entonox®)	96
Oxygen	97
Paracetamol	98
Salbutamol	99

APPENDIX 1

MEDICATION FORMULARY

CLINICAL LEVEL:



Medication	Aspirin
Class	Platelet aggregation inhibitor
Descriptions	Anti-inflammatory agent and an inhibitor of platelet function. Useful agent in the treatment of various thromboembolic diseases such as acute myocardial infarction.
Presentation	300 mg dispersible tablet
Administration	Orally (PO) – dispersed in water, or to be chewed – if not dispersible form (CPG: 5/6.4.10, 4.4.10, 1/2/3.4.10)
Indications	Cardiac chest pain or suspected Myocardial Infarction
Contraindications	Active symptomatic gastrointestinal (GI) ulcer Bleeding disorder (e.g. haemophilia) Known severe adverse reaction Patients < 16 years old
Usual Dosages	Adult: 300 mg tablet Paediatric: Contraindicated
Pharmacology/Action	Antithrombotic Inhibits the formation of thromboxane A ₂ , which stimulates platelet aggregation and artery constriction. This reduces clot/thrombus formation in an MI.
Side effects	Epigastric pain and discomfort Bronchospasm Gastrointestinal haemorrhage
Long-term effects	Generally mild and infrequent but incidence of gastro-intestinal irritation with slight asymptomatic blood loss, increased bleeding time, bronchospasm and skin reaction in hypersensitive patients.
Additional information	Aspirin 300 mg is indicated for cardiac chest pain regardless if patient is on anticoagulants or is already on aspirin. If the patient has swallowed an aspirin (enteric coated) preparation without chewing it, the patient should be regarded as not having taken any aspirin; administer 300 mg PO.

APPENDIX 1

MEDICATION FORMULARY

CLINICAL LEVEL:    

Medication	Epinephrine (1:1,000)												
Class	Sympathetic agonist												
Description	Naturally occurring catecholamine. It is a potent alpha and beta adrenergic stimulant; however, its effect on beta receptors is more profound.												
Presentation	Pre-filled syringe, ampoule or Auto injector (for EMT use) 1 mg/1 mL (1:1,000)												
Administration	Intramuscular (IM) (CPG: 5/6.4.15, 4.4.15, 2/3.4.16, 5/6.7.31, 4.7.31, 2/3.7.31)												
Indications	Severe anaphylaxis												
Contraindications	None known												
Usual Dosages	<p>Adult: 0.5 mg (500 mcg) IM (0.5 mL of 1: 1,000) EMT & (EFR assist patient) 0.3 mg (Auto injector) Repeat every 5 minutes prn</p> <p>Paediatric:</p> <table border="0"> <tr> <td>< 6 months:</td> <td>0.05 mg (50 mcg) IM (0.05 mL of 1:1 000)</td> </tr> <tr> <td>6 months to 5 years:</td> <td>0.125 mg (125 mcg) IM (0.13 mL of 1:1 000)</td> </tr> <tr> <td>6 to 8 years:</td> <td>0.25 mg (250 mcg) IM (0.25 mL of 1:1 000)</td> </tr> <tr> <td>> 8 years:</td> <td>0.5 mg (500 mcg) IM (0.5 mL of 1:1 000)</td> </tr> </table> <p>EMT & (EFR assist patient):</p> <table border="0"> <tr> <td>6 months < 10 years:</td> <td>0.15 mg (Auto injector)</td> </tr> <tr> <td>≥ 10 years:</td> <td>0.3 mg (Auto injector)</td> </tr> </table> Repeat every 5 minutes prn	< 6 months:	0.05 mg (50 mcg) IM (0.05 mL of 1:1 000)	6 months to 5 years:	0.125 mg (125 mcg) IM (0.13 mL of 1:1 000)	6 to 8 years:	0.25 mg (250 mcg) IM (0.25 mL of 1:1 000)	> 8 years:	0.5 mg (500 mcg) IM (0.5 mL of 1:1 000)	6 months < 10 years:	0.15 mg (Auto injector)	≥ 10 years:	0.3 mg (Auto injector)
< 6 months:	0.05 mg (50 mcg) IM (0.05 mL of 1:1 000)												
6 months to 5 years:	0.125 mg (125 mcg) IM (0.13 mL of 1:1 000)												
6 to 8 years:	0.25 mg (250 mcg) IM (0.25 mL of 1:1 000)												
> 8 years:	0.5 mg (500 mcg) IM (0.5 mL of 1:1 000)												
6 months < 10 years:	0.15 mg (Auto injector)												
≥ 10 years:	0.3 mg (Auto injector)												
Pharmacology/Action	Alpha and beta adrenergic stimulant Reversal of laryngeal oedema & bronchospasm in anaphylaxis Antagonises the effects of histamine												
Side effects	Palpitations Tachyarrhythmias Hypertension Angina-like symptoms												
Additional information	N.B. Double check the concentration on pack before use												

APPENDIX 1 MEDICATION FORMULARY

CLINICAL LEVEL:   

Medication	Glucagon
Class	Hormone and Antihypoglycaemic
Description	Glucagon is a protein secreted by the alpha cells of the Islets of Langerhans in the pancreas. It is used to increase the blood glucose level in cases of hypoglycaemia in which an IV cannot be immediately placed.
Presentation	1 mg vial powder and solution for reconstitution (1 mL)
Administration	Intramuscular (IM) (CPG: 5/6.4.19, 4.4.19, 5/6.7.32, 4.7.32)
Indications	Hypoglycaemia in patients unable to take oral glucose or unable to gain IV access, with a blood glucose level < 4 mmol/L
Contraindications	Known severe adverse reaction Pheochromocytoma
Usual Dosages	Adult: 1 mg IM Paediatric: ≤ 8 years 0.5 mg (500 mcg) IM > 8 years 1 mg IM
Pharmacology/Action	Glycogenolysis Increases plasma glucose by mobilising glycogen stored in the liver
Side effects	Rare, may cause hypotension, dizziness, headache, nausea & vomiting
Additional information	May be ineffective in patients with low stored glycogen e.g. prior use in previous 24 hours, alcoholic patients with liver disease. Store in refrigerator Protect from light

APPENDIX 1 MEDICATION FORMULARY

CLINICAL LEVEL:



Medication	Glucose gel
Class	Antihypoglycaemic
Description	Synthetic glucose paste
Presentation	Glucose gel in a tube or sachet
Administration	Buccal administration: Administer gel to the inside of the patient's cheek and gently massage the outside of the cheek. (CPG: 5/6.4.19, 4.4.19, 2/3.4.19, 5/6.7.32, 4.7.32)
Indications	Hypoglycaemia Blood glucose < 4 mmol/L EFR – Known diabetic with confusion or altered levels of consciousness
Contraindications	Known severe adverse reaction
Usual Dosages	Adult: 10 – 20 g buccal Repeat prn Paediatric: ≤ 8 years; 5 – 10 g buccal >8 years: 10 – 20 g buccal Repeat prn
Pharmacology/Action	Increases blood glucose levels
Side effects	May cause vomiting in patients under the age of five if administered too quickly
Additional information	Glucose gel will maintain glucose levels once raised but should be used secondary to Dextrose to reverse hypoglycaemia. Proceed with caution: Patients with airway compromise Altered level of consciousness

APPENDIX 1 MEDICATION FORMULARY

CLINICAL LEVEL:    

Medication	Glyceryl Trinitrate (GTN)
Class	Nitrate
Description	Special preparation of Glyceryl trinitrate in an aerosol form that delivers precisely 0.4 mg of Glyceryl trinitrate per spray.
Presentation	Aerosol spray: metered dose 0.4 mg (400 mcg)
Administration	Sublingual (SL): Hold the pump spray vertically with the valve head uppermost Place as close to the mouth as possible and spray under the tongue The mouth should be closed after each dose (CPG: 5/6.3.5, 4.4.10, 5/6.4.10)
Indications	Angina Suspected Myocardial Infarction (MI) EFRs may assist with administration Advanced Paramedic and Paramedic - Pulmonary oedema
Contraindications	SBP < 90 mmHg Viagra or other phosphodiesterase type 5 inhibitors (Sildenafil, Tadalafil and Vardenafil) used within previous 24 hours. Known severe adverse reaction.
Usual Dosages	Adult: Angina or MI: 0.4 mg (400 mcg) Sublingual Repeat at 3-5 min intervals, Max: 1.2 mg (EFRs 0.4 mg sublingual max, assist patient) Pulmonary oedema; 0.8 mg (800 mcg) sublingual Repeat x 1 Paediatric: Not indicated
Pharmacology/Action	Vasodilator Releases nitric oxide which acts as a vasodilator. Dilates coronary arteries particularly if in spasm increasing blood flow to myocardium. Dilates systemic veins reducing venous return to the heart (pre load) and thus reduces the heart's workload. Reduces BP
Side effects	Headache Transient Hypotension Flushing Dizziness
Additional information	If the pump is new or has not been used for a week or more, the first spray should be released into the air.

APPENDIX 1
MEDICATION FORMULARY

CLINICAL LEVEL:   

Medication	Ibuprofen
Class	Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)
Description	It is an anti-inflammatory analgesic
Presentation	Suspension 100 mg in 5 mL 200 mg tablet, 400 mg tablet
Administration	Orally (PO) (CPG: 4/5/6.2.6, 4/5/6.7.5)
Indications	Mild to moderate pain
Contraindications	Not suitable for children under 3 months Patient with history of asthma exacerbated by aspirin Pregnancy Peptic ulcer disease Known severe adverse reaction
Usual Dosages	Adult: 400 mg PO Paediatric: 10 mg/Kg PO
Pharmacology/Action	Suppresses prostaglandins, which cause pain via the inhibition of cyclooxygenase (COX). Prostaglandins are released by cell damage and inflammation.
Side effects	Skin rashes, gastrointestinal intolerance and bleeding
Long-term side effects	Occasionally gastrointestinal bleeding and ulceration occurs May also cause acute renal failure, interstitial nephritis and NSAID-associated nephropathy
Additional information	If Ibuprofen administered in previous 6 hours, adjust the dose downward by the amount given by other sources resulting in a maximum of 10 mg/Kg. Caution with significant burns or poor perfusion due to risk of kidney failure. Caution if concurrent NSAIDs use.

APPENDIX 1 MEDICATION FORMULARY

CLINICAL LEVEL:   

Medication	Naloxone
Class	Narcotic antagonist
Description	Effective in management and reversal of overdoses caused by narcotics or synthetic narcotic agents.
Presentation	Ampoules 0.4 mg in 1 mL (400 mcg /1 mL) or pre-loaded syringe
Administration	Intravenous (IV) Intramuscular (IM) Subcutaneous (SC) Intraosseous (IO) Intranasal (IN) (CPG: 6.4.22, 4/5.4.22, 5/6.5.2, 4/5/6.7.11)
Indications	Inadequate respiration and/or ALoC following known or suspected narcotic overdose
Contraindications	Known severe adverse reaction
Usual Dosages	<p>Adult: 0.4 mg (400 mcg) IV/IO (AP) 0.4 mg (400 mcg) IM or SC (P) 0.8 mg (800 mcg) IN (EMT) Repeat after 3 min prn to a Max 2 mg</p> <p>Paediatric: 0.01 mg/Kg (10 mcg/Kg) IV/IO (AP) 0.01 mg/Kg (10 mcg/Kg) IM/SC (P) 0.02 mg/Kg (20 mcg/Kg) IN (EMT) Repeat dose prn to maintain opioid reversal to Max 0.1 mg/Kg or 2 mg</p>
Pharmacology/Action	Narcotic antagonist Reverse the respiratory depression and analgesic effect of narcotics
Side effects	Acute reversal of narcotic effect ranging from nausea & vomiting to agitation and seizures
Additional information	Use with caution in pregnancy. Administer with caution to patients who have taken large dose of narcotics or are physically dependent. Rapid reversal will precipitate acute withdrawal syndrome. Prepare to deal with aggressive patients.

APPENDIX 1 MEDICATION FORMULARY

CLINICAL LEVEL:   

Medication	Nitrous Oxide 50% and Oxygen 50% (Entonox®)
Class	Analgesic
Description	Potent analgesic gas contains a mixture of both nitrous oxide and oxygen
Presentation	Cylinder, coloured blue with white and blue triangles on cylinder shoulders Medical gas: 50% Nitrous Oxide Et 50% Oxygen
Administration	Self-administered Inhalation by demand valve with face-mask or mouthpiece (CPG: 4/5/6.2.6, 5/6.5.1, 4.5.1, 5/6.5.6, 4/5/6.7.5)
Indications	Pain relief
Contraindications	Altered level of consciousness Chest Injury/Pneumothorax Shock Recent scuba dive Decompression sickness Intestinal obstruction Inhalation Injury Carbon monoxide (CO) poisoning Known severe adverse reaction
Usual Dosages	Adult: Self-administered until pain relieved Paediatric: Self-administered until pain relieved
Pharmacology/Action	Analgesic agent gas: - CNS depressant - Pain relief
Side effects	Disinhibition Decreased level of consciousness Light-headedness
Additional information	Do not use if patient unable to understand instructions. In cold temperatures warm cylinder and invert to ensure mix of gases. Advanced Paramedics may use discretion with minor chest injuries. Brand name: Entonox®. Has an addictive property. Caution when using Entonox for greater than one hour for Sickle Cell Crisis.

APPENDIX 1 MEDICATION FORMULARY

CLINICAL LEVEL:



Medication	Oxygen
Class	Gas
Description	Odourless, tasteless, colourless gas necessary for life
Presentation	D, E or F cylinders, coloured black with white shoulders CD cylinder; white cylinder Medical gas
Administration	Inhalation via: High concentration reservoir (non-rebreather) mask Simple face mask Venturi mask Tracheostomy mask Nasal cannulae Bag Valve Mask (CPG: Oxygen is used extensively throughout the CPGs)
Indications	Absent/inadequate ventilation following an acute medical or traumatic event SpO ₂ < 94% adults and < 96% paediatrics SpO ₂ < 92% for patients with acute exacerbation of COPD
Contraindications	Bleomycin lung injury
Usual Dosages	<p>Adult: Cardiac and respiratory arrest or Sickle Cell Crisis; 100% Life threats identified during primary survey; 100% until a reliable SpO₂ measurement obtained then titrate O₂ to achieve SpO₂ of 94% - 98% For patients with acute exacerbation of COPD, administer O₂ titrate to achieve SpO₂ 92% or as specified on COPD Oxygen Alert Card All other acute medical and trauma titrate O₂ to achieve SpO₂ 94% -98%</p> <p>Paediatric: Cardiac and respiratory arrest or Sickle Cell Crisis; 100% Life threats identified during primary survey; 100% until a reliable SpO₂ measurement obtained then titrate O₂ to achieve SpO₂ of 96% - 98% All other acute medical and trauma titrate O₂ to achieve SpO₂ of 96% - 98%</p>
Pharmacology/Action	Oxygenation of tissue/organs
Side effects	Prolonged use of O ₂ with chronic COPD patients may lead to reduction in ventilation stimulus.
Additional information	A written record must be made of what oxygen therapy is given to every patient. Documentation recording oximetry measurements should state whether the patient is breathing air or a specified dose of supplemental oxygen. Consider humidifier if oxygen therapy for paediatric patients is > 30 minute duration. Caution with paraquat poisoning, administer oxygen if SpO ₂ < 92% Avoid naked flames, powerful oxidising agent.

APPENDIX 1
MEDICATION FORMULARY

CLINICAL LEVEL:   

Medication	Paracetamol
Class	Analgesic and antipyretic
Description	Paracetamol is used to reduce pain and body temperature
Presentation	Rectal suppository 180 mg, 90 mg and 60 mg Suspension 120 mg in 5 mL or 250 mg in 5 mL 500 mg tablet
Administration	Per Rectum (PR) Orally (PO) (CPG: 4/5/6.2.6, 4/5/6.4.24, 4/5/6.7.5, 4/5/6.7.35)
Indications	Pyrexia Minor or moderate pain (1 - 6 on pain scale) for adult and paediatric patients
Contraindications	Known severe adverse reaction Chronic liver disease < 1 month old
Usual Dosages	Adult: 1 g PO Paediatric: PR (AP) PO (AP, P & EMT) > 1 mth < 1 year - 90 mg PR 20 mg/Kg PO 1-3 years - 180 mg PR 4-8 years - 360 mg PR
Pharmacology/Action	Analgesic – central prostaglandin inhibitor Antipyretic – prevents the hypothalamus from synthesising prostaglandin E, inhibiting the body temperature from rising further
Side effects	None
Long-term side effects	Long-term use at high dosage or over dosage can cause liver damage and less frequently renal damage
Additional information	Note: Paracetamol is contained in Paracetamol Suspension and other over the counter drugs. Consult with parent/guardian in relation to medication prior to arrival on scene. For PR use be aware of modesty of patient, should be administered in presence of a 2 nd person. If Paracetamol administered in previous 4 hours, adjust the dose downward by the amount given by other sources resulting in a maximum of 20 mg/Kg.

APPENDIX 1
MEDICATION FORMULARY

CLINICAL LEVEL:    

Medication	Salbutamol
Class	Sympathetic agonist
Description	Sympathomimetic that is selective for beta-2 adrenergic receptors
Presentation	Nebule 2.5 mg in 2.5 mL Nebule 5 mg in 2.5 mL Aerosol inhaler: metered dose 0.1 mg (100 mcg)
Administration	Nebuliser (NEB) Inhalation via aerosol inhaler (CPG: 4/5/6.3.3, 4/5/6.3.4, 3.3.4, 5/6.4.15, 4.4.15, 2/3.4.16, 4/5/6.6.10, 4/5/6.7.12, 3.7.12, 5/6.7.31, 4.7.31, 2/3.7.31)
Indications	Bronchospasm Exacerbation of COPD Respiratory distress following submersion incident
Contraindications	Known severe adverse reaction
Usual Dosages	Adult: 5 mg NEB (or 0.1 mg metered aerosol spray x 5) Repeat at 5 min prn (EFRs: 0.1 mg metered aerosol spray x 5, assist patient) Paediatric: < 5 yrs - 2.5 mg NEB (or 0.1 mg metered aerosol spray x 3) ≥ 5 yrs - 5 mg NEB (or 0.1 mg metered aerosol spray x 5) Repeat at 5 min prn (EFRs: 0.1 mg metered aerosol spray x 2, assist patient)
Pharmacology/Action	Beta-2 agonist Bronchodilation Relaxation of smooth muscle
Side effects	Tachycardia. Tremors Tachyarrhythmias High doses may cause hypokalaemia
Additional information	It is more efficient to use a volumizer in conjunction with an aerosol inhaler when administering Salbutamol. If an oxygen driven nebuliser is used to administer Salbutamol for a patient with acute exacerbation of COPD it should be limited to 6 minutes maximum.

APPENDIX 2 MEDICATIONS & SKILLS MATRIX

NEW FOR 2014

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Burns care			✓	✓	✓	✓	✓
Soft tissue injury			✓	✓	✓	✓	✓
SpO ₂ monitoring				✓			
Move and secure a patient to a paediatric board					✓		
Ibuprofen PO					✓		
Salbutamol Nebule					✓		
Subcutaneous injection					✓	✓	
Naloxone IN					✓	✓	✓
Pain assessment					✓	✓	✓
Haemostatic agent					✓	✓	✓
End Tidal CO ₂ monitoring						✓	
Hydrocortisone IM						✓	
Ipratropium Bromide Nebule						✓	
CPAP / BiPAP						✓	✓
Naloxone SC						✓	✓
Nasal pack						✓	✓
Ticagrelor						✓	✓
Treat and referral						✓	✓
Tranexamic Acid							✓

CARE MANAGEMENT INCLUDING THE ADMINISTRATION OF MEDICATIONS AS PER LEVEL OF TRAINING AND DIVISION ON THE PHECC REGISTER AND RESPONDER LEVELS.

Pre-Hospital responders and practitioners shall only provide care management including medication administration for which they have received specific training. Practitioners must be privileged by a licensed CPG provider to administer specific medications and perform specific clinical interventions.

KEY		
✓	=	Authorised under PHECC CPGs
URMPIO	=	Authorised under PHECC CPGs under registered medical practitioner's instructions only
APO	=	Authorised under PHECC CPGs to assist practitioners only (when applied to EMT, to assist Paramedic or higher clinical levels)
SA	=	Authorised subject to special authorisation as per CPG
BTEC	=	Authorised subject to Basic Tactical Emergency Care rules

APPENDIX 2
MEDICATIONS & SKILLS MATRIX

MEDICATIONS

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Aspirin PO	✓	✓	✓	✓	✓	✓	✓
Oxygen		✓		✓	✓	✓	✓
Glucose Gel Buccal				✓	✓	✓	✓
GTN SL				✓SA	✓	✓	✓
Salbutamol Aerosol				✓SA	✓	✓	✓
Epinephrine (1:1,000) auto injector				✓SA	✓	✓	✓
Glucagon IM					✓	✓	✓
Nitrous oxide & Oxygen (Entonox®)					✓	✓	✓
Naloxone IN					✓	✓	✓
Paracetamol PO					✓	✓	✓
Ibuprofen PO					✓	✓	✓
Salbutamol nebule					✓	✓	✓
Morphine IM					URMPIO	URMPIO	✓SA
Clopidogrel PO						✓	✓
Epinephrine (1: 1,000) IM						✓	✓
Hydrocortisone IM						✓	✓
Ipratropium Bromide Nebule						✓	✓
Midazolam IM/Buccal/IN						✓	✓
Naloxone IM/SC						✓	✓
Ticagrelor						✓	✓
Dextrose 10% IV						✓SA	✓
Hartmann's Solution IV/IO						✓SA	✓
Sodium Chloride 0.9% IV/IO						✓SA	✓
Amiodarone IV/IO							✓
Atropine IV/IO							✓
Benzympenicillin IM/IV/IO							✓
Cyclizine IV							✓
Diazepam IV/PR							✓
Epinephrine (1:10,000) IV/IO							✓
Fentanyl IN							✓
Furosemide IV/IM							✓
Hydrocortisone IV							✓
Lorazepam PO							✓
Magnesium Sulphate IV							✓
Midazolam IV							✓

APPENDIX 2 MEDICATIONS & SKILLS MATRIX

MEDICATIONS (contd)

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Morphine IV/PO							✓
Naloxone IV/IO							✓
Nifedipine PO							✓
Ondansetron IV							✓
Paracetamol PR							✓
Sodium Bicarbonate IV/ IO							✓
Syntometrine IM							✓
Tranexamic Acid							✓
Enoxaparin IV/SC							✓ SA
Lidocaine IV							✓ SA
Tenecteplase IV							✓ SA

AIRWAY & BREATHING MANAGEMENT

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
FBAO management	✓	✓	✓	✓	✓	✓	✓
Head tilt chin lift	✓	✓	✓	✓	✓	✓	✓
Pocket mask	✓	✓	✓	✓	✓	✓	✓
Recovery position	✓	✓	✓	✓	✓	✓	✓
Non rebreather mask		✓		✓	✓	✓	✓
OPA		✓		✓	✓	✓	✓
Suctioning		✓		✓	✓	✓	✓
Venturi mask		✓		✓	✓	✓	✓
SpO ₂ monitoring		✓ SA		✓	✓	✓	✓
Jaw Thrust				✓	✓	✓	✓
Nasal cannula		✓		✓	✓	✓	✓
BVM		✓		✓ SA	✓	✓	✓
NPA				BTEC	BTEC	✓	✓
Supraglottic airway adult (uncuffed)		✓			✓	✓	✓
Oxygen humidification					✓	✓	✓
Supraglottic airway adult (cuffed)					✓ SA	✓	✓
CPAP / BiPAP						✓	✓
Non-invasive ventilation device						✓	✓
Peak Expiratory Flow						✓	✓

APPENDIX 2 MEDICATIONS & SKILLS MATRIX

AIRWAY & BREATHING MANAGEMENT *(contd)*

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
End Tidal CO ₂ monitoring						✓	✓
Supraglottic airway paediatric						✓ SA	✓
Endotracheal intubation							✓
Laryngoscopy and Magill forceps							✓
Needle cricothyrotomy							✓
Needle thoracocentesis							✓

CARDIAC

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
AED adult & paediatric	✓	✓	✓	✓	✓	✓	✓
CPR adult, child & infant	✓	✓	✓	✓	✓	✓	✓
Recognise death and resuscitation not indicated	✓	✓	✓	✓	✓	✓	✓
Targeted temperature management		✓ SA			✓	✓	✓
CPR newly born					✓	✓	✓
ECG monitoring (lead II)					✓	✓	✓
Mechanical assist CPR device					✓	✓	✓
12 lead ECG						✓	✓
Cease resuscitation - adult						✓	✓
Manual defibrillation						✓	✓

HAEMORRHAGE CONTROL

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Direct pressure			✓	✓	✓	✓	✓
Nose bleed			✓	✓	✓	✓	✓
Haemostatic agent					✓	✓	✓
Tourniquet use				BTEC	BTEC	✓	✓
Nasal pack						✓	✓
Pressure points						✓	✓

APPENDIX 2 MEDICATIONS & SKILLS MATRIX

MEDICATION ADMINISTRATION

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Oral	✓	✓	✓	✓	✓	✓	✓
Buccal route				✓	✓	✓	✓
Per aerosol (inhaler) + spacer				✓SA	✓	✓	✓
Sublingual				✓SA	✓	✓	✓
Intramuscular injection					✓	✓	✓
Intranasal					✓	✓	✓
Per nebuliser					✓	✓	✓
Subcutaneous injection					✓	✓	✓
IV & IO Infusion maintenance						✓SA	✓
Infusion calculations							✓
Intraosseous injection/infusion							✓
Intravenous injection/infusion							✓
Per rectum							✓

TRAUMA

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Burns care			✓	✓	✓	✓	✓
Cervical spine manual stabilisation			✓	✓	✓	✓	✓
Application of a sling			✓	✓	✓	✓	✓
Soft tissue injury			✓	✓	✓	✓	✓
Cervical collar application				✓	✓	✓	✓
Helmet stabilisation/removal				✓	✓	✓	✓
Splinting device application to upper limb				✓	✓	✓	✓
Move and secure patient to a long board				✓SA	✓	✓	✓
Rapid Extraction				✓SA	✓	✓	✓
Log roll				APO	✓	✓	✓
Move patient with a carrying sheet				APO	✓	✓	✓
Move patient with an orthopaedic stretcher				APO	✓	✓	✓
Splinting device application to lower limb				APO	✓	✓	✓
Secure and move a patient with an extrication device				APO	APO	✓	✓

APPENDIX 2
MEDICATIONS & SKILLS MATRIX

TRAUMA (contd)

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Pelvic Splinting device				BTEC	✓	✓	✓
Move and secure patient into a vacuum mattress				BTEC	✓	✓	✓
Active re-warming					✓	✓	✓
Move and secure a patient to a paediatric board					✓	✓	✓
Traction splint application					APO	✓	✓
Spinal Injury Decision						✓	✓
Taser gun barb removal						✓	✓
Reduction dislocated patella							✓

OTHER

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Assist in the normal delivery of a baby				APO	✓	✓	✓
De-escalation and breakaway skills					✓	✓	✓
Glucometry					✓	✓	✓
Broselow tape						✓	✓
Delivery Complications						✓	✓
External massage of uterus						✓	✓
Intraosseous cannulation							✓
Intravenous cannulation							✓
Urinary catheterisation							✓

PATIENT ASSESSMENT

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Assess responsiveness	✓	✓	✓	✓	✓	✓	✓
Check breathing	✓	✓	✓	✓	✓	✓	✓
FAST assessment	✓	✓	✓	✓	✓	✓	✓
Capillary refill			✓	✓	✓	✓	✓
AVPU			✓	✓	✓	✓	✓
Breathing & pulse rate			✓	✓	✓	✓	✓

APPENDIX 2
MEDICATIONS & SKILLS MATRIX

PATIENT ASSESSMENT (contd)

CLINICAL LEVEL	CFR-C	CFR-A	FAR/OFA	EFR	EMT	P	AP
Primary survey			✓	✓	✓	✓	✓
SAMPLE history			✓	✓	✓	✓	✓
Secondary survey			✓	✓	✓	✓	✓
CSM assessment				✓	✓	✓	✓
Rule of Nines				✓	✓	✓	✓
Assess pupils				✓	✓	✓	✓
Blood pressure				✓SA	✓	✓	✓
Capacity evaluation					✓	✓	✓
Do Not Attempt Resuscitation					✓	✓	✓
Paediatric Assessment Triangle					✓	✓	✓
Pain assessment					✓	✓	✓
Patient Clinical Status					✓	✓	✓
Pre-hospital Early Warning Score					✓	✓	✓
Pulse check (cardiac arrest)		✓SA			✓	✓	✓
Temperature °C					✓	✓	✓
Triage sieve					✓	✓	✓
Chest auscultation					✓	✓	✓
GCS					✓	✓	✓
Treat and referral					✓	✓	✓
Triage sort					✓	✓	✓

APPENDIX 3 CRITICAL INCIDENT STRESS MANAGEMENT

Your Psychological Well-Being

As a Practitioner it is extremely important for your psychological well-being that you do not expect to save every critically ill or injured patient that you treat. For a patient who is not in hospital, whether they survive a cardiac arrest or multiple trauma depends on a number of factors including any other medical condition the patient has. Your aim should be to perform your interventions well and to administer the appropriate medications within your scope of practice. However sometimes you may encounter a situation which is highly stressful for you, giving rise to Critical Incident Stress (CIS). A critical incident is an incident or event which may overwhelm or threaten to overwhelm our normal coping responses. As a result of this we can experience CIS.

SYMPTOMS OF CIS INCLUDE SOME OR ALL OF THE FOLLOWING:

Examples of physical symptoms:

- Feeling hot and flushed, sweating a lot
- Dry mouth, churning stomach
- Diarrhoea and digestive problems
- Needing to urinate often
- Muscle tension
- Restlessness, tiredness, sleep difficulties, headaches
- Increased drinking or smoking
- Overeating, or loss of appetite
- Loss of interest in sex
- Racing heart, breathlessness and rapid breathing

Examples of psychological symptoms:

- Feeling overwhelmed
- Loss of motivation
- Dreading going to work
- Becoming withdrawn
- Racing thoughts
- Confusion
- Not looking after yourself properly
- Difficulty making decisions
- Poor concentration
- Poor memory
- Anger
- Anxiety
- Depression

Post-Traumatic Stress Reactions

Normally the symptoms of Critical Incident Stress subside within a few weeks or less. Sometimes however, they may persist and develop into a post-traumatic stress reaction and you may also experience emotional reactions.

Anger at the injustice and senselessness of it all.

Sadness and depression caused by an awareness of how little can be done for people who are severely injured and dying, sense of a shortened future, poor concentration, not being able to remember things as well as before.

Guilt caused by believing that you should have been able to do more or that you could have acted differently.

Fear of 'breaking down' or 'losing control', not having done all you could have done, being blamed for something or a similar event happening to you or your loved ones.

APPENDIX 3 CRITICAL INCIDENT STRESS MANAGEMENT

Avoiding the scene of the trauma or anything that reminds you of it.

Intrusive thoughts in the form of memories or flashbacks which cause distress and the same emotions as you felt at the time.

Irritability outbursts of anger, being easily startled and constantly being on guard for threats.

Feeling numb leading to a loss of your normal range of feelings, for example, being unable to show affection, feeling detached from others.

EXPERIENCING SIGNS OF EXCESSIVE STRESS

If the range of physical, emotional and behavioural signs and symptoms already mentioned do not reduce over time (for example, after two weeks), it is important that you get support and help.

Where to find help?

Your own CPG approved organisation will have a CISM support network or system.

We recommend that you contact them for help and advice. (i.e. your peer support worker/coordinator/staff support officer).

- For a self-help guide, please go to www.cismnetworkireland.ie
- The NAS CISM/ CISM Network published a booklet called 'Critical Incident Stress Management for Emergency Personnel'. It can be purchased by emailing info@cismnetworkireland.ie
- The NAS CISM committee in partnership with PHECC developed an eLearning CISM Stress Awareness Training (SAT) module. It can be accessed by all PHECC registered practitioners using their PHECC eLearning username and password. In due course PHECC will launch a CISM SAT module for non-PHECC registered personnel.
- See a health professional who specialises in traumatic stress.

APPENDIX 4 CPG UPDATES FOR EMERGENCY MEDICAL TECHNICIANS

CPG updates 2014

For administrative purposes the numbering system on some CPGs has been changed.

The paediatric age range has been extended to reflect the new national paediatric age (≤ 15 years), as outlined by National Clinical Programme for Paediatrics and Neonatology.

CPGs that have content changes are outlined below.

Updated CPGs from the 2012 version.

CPGs	The principal differences are	Theory	Skills
CPG 4/5/6.2.1 Primary Survey Medical – Adult	EMTs, who have completed the BTEC course, may be privileged by a licenced CPG provider to insert an NPA following appropriate training	✓	BTEC only
CPG 4/5/6.2.2 Primary Survey Trauma – Adult	EMTs, who have completed the BTEC course, may be privileged by a licenced CPG provider to insert an NPA following appropriate training	✓	BTEC only
CPG 4/5/6.2.6 Pain Management – Adult	Delete 'Minor pain (2 to 3 on pain scale)' replace with 'Mild pain (1 to 3 on pain scale)'	✓	x
	Change Moderate pain to '4 to 6 on the pain scale'	✓	x
	Change Severe pain to '≥ 7 on the pain scale'	✓	x
	Add Fentanyl IN for advanced paramedic practice	✓	x
	Add Ibuprofen PO for EMT practice	✓	✓
CPG 4.3.1 Advanced Airway Management – Adult	Special authorisation may be given to EMTs to insert a cuffed supraglottic airway subject to maintaining competence and Medical Director authorisation	✓	✓ if authorised
CPG 4/5/6.3.2 Inadequate Ventilations – Adult	This CPG replaces Inadequate Respirations – Adult (5/6.3.2 and 4.3.2) incorporating all three practitioner levels in one CPG	✓	x
	This CPG outlines generic care for all patients with inadequate ventilation and then offers pathways for specific clinical issues	✓	x

APPENDIX 4 CPG UPDATES FOR EMERGENCY MEDICAL TECHNICIANS

CPGs	The principal differences are	Theory	Skills
CPG 4/5/6.3.3 Exacerbation of COPD	This CPG incorporating all three practitioner levels in one CPG replacing 4.3.3 at EMT level Peak expiratory flow measurement is now within the scope of practice for paramedics Salbutamol Neb is now within the scope of practice for EMTs Ipratropium bromide Neb is now within the scope of practice for paramedics	✓ ✓ ✓ ✓	x x ✓ x
CPG 4/5/6.4.11 Symptomatic Bradycardia – Adult	The dose of Atropine has been increased from 0.5 mg to 0.6 mg Add 'NaCl infusion 250 mL (repeat by one)' Insert information box; 'Titrate Atropine to effect (HR > 60)'	✓ ✓ ✓	x x x
CPG 4.4.15 Allergic Reaction/ Anaphylaxis – Adult	Salbutamol NEB is now within the scope of practice for EMTs The conditions for use of Epinephrine auto injector has been changed; it is now indicated for all patients with severe anaphylaxis regardless of whether it has been previously prescribed or not.	✓ ✓	x x
CPG 4/5/6.4.17 Epistaxis	Digital pressure has been increased to 15 minutes The insertion of a proprietary nasal pack is now within the scope of practice for paramedics and advanced paramedics	✓ ✓	x x
CPG 4.4.21 Hypothermia	Paramedic has been removed from this CPG Warmed O2 has been removed	✓ ✓	x x
CPG 4/5.4.22 Poisons – Adult	The methods of introduction of a poison have been removed Naloxone has been added to this CPG for opiate induced poison Naloxone IN is now within the scope of practice for EMTs and paramedics The absolute contraindication for O2 has been removed following paraquat poisoning	✓ ✓ ✓ ✓	x x ✓ x
CPG 4/5/6.4.24 Sepsis – Adult	This CPG replaces Septic Shock – Adult It authorises the administration of Paracetamol for pyrexia patients It authorises the administration, by advanced paramedics, of Benzylpenicillin for severe sepsis. Advanced paramedics may consider additional aliquots of NaCl to maintain systolic BP > 100 mmHg	✓ ✓ ✓ ✓	x x x x

APPENDIX 4 CPG UPDATES FOR EMERGENCY MEDICAL TECHNICIANS

CPGs	The principal differences are	Theory	Skills
CPG 4/5/6.6.1 Burns – Adult	Add 'Caution with hypothermia'	✓	x
CPG 4/5/6.6.3 External Haemorrhage – Adult	This CPG has been updated to reflect the importance of managing catastrophic haemorrhage immediately. Dressings impregnated with haemostatic agents are now within the scope of practice for EMTs, paramedics and advanced paramedics. EMTs, who have completed the BTEC course, may be privileged by a licenced CPG provider to apply a tourniquet.	✓ ✓ ✓	x ✓ BTEC only
CPG 4.6.5 Head Injury – Adult	Add V as a rationale for requesting ALS Add 'consider mechanism of injury; is spinal immobilisation indicated?' Replace 'apply cervical collar' and 'secure to long board' with 'immobilise spine appropriately'	✓ ✓ ✓	x x x
CPG 4/5/6.6.7 Limb Injury – Adult	Fractured neck of femur has been included With a fractured neck of femur, if the transport time to ED is > 20 minutes, ALS should be requested. With a fractured neck of femur advanced paramedics should consider NaCl infusion	✓ ✓ ✓	x x x
CPG 4.6.8 Shock from Blood Loss – Adult	The signs of poor perfusion have been presented in an ABCDE format	✓	x
CPG 4/5/6.6.10 Submersion Incident	Salbutamol is now within the scope of practice for EMTs	✓	✓
CPG 4/5/6.7.4 Secondary Survey – Paediatric	The estimated weight formula has been updated; Neonate = 3.5 Kg Six months = 6 Kg One to five years = (age x 2) + 8 Kg Greater than 5 years = (age x 3) + 7 Kg	✓	x

APPENDIX 4 CPG UPDATES FOR EMERGENCY MEDICAL TECHNICIANS

CPGs	The principal differences are	Theory	Skills
CPG 4/5/6.7.5 Pain Management – Paediatric	<p>Pain assessment recommendations; < 5 years use FLACC scale 5 – 7 years use Wong Baker scale ≥ 8 years use analogue pain scale</p> <p>Delete 'Minor pain (2 to 3 on pain scale)' replace with 'Mild pain (1 to 3 on pain scale)'</p> <p>Change Moderate pain to '4 to 6 on the pain scale'</p> <p>Change Severe pain to '≥ 7 on the pain scale'</p> <p>Fentanyl IN is now within the scope of practice for advanced paramedics</p> <p>Ibuprofen PO is now within the scope of practice for EMTs</p>	<p>✓</p> <p>✓</p> <p>✓</p> <p>✓</p> <p>✓</p> <p>✓</p>	<p>✓</p> <p>x</p> <p>x</p> <p>x</p> <p>x</p> <p>✓</p>
CPG 4/5/6.7.11 Inadequate Ventilations – Paediatric	<p>This CPG replaces Inadequate Respirations – Paediatric (5/6.7.5 and 4.7.5) incorporating all three practitioner levels in one CPG</p> <p>This CPG outlines generic care for all patients with inadequate ventilation and then offers pathways for specific clinical issues</p> <p>Naloxone IN is now within the scope of practice for EMTs, paramedics and advanced paramedics.</p>	<p>✓</p> <p>✓</p> <p>✓</p>	<p>x</p> <p>x</p> <p>✓</p>
CPG 4/5/6.7.24 Symptomatic Bradycardia – Paediatric	<p>The routine ventilations has been changed to ventilations if hypoxic.</p> <p>Unresponsive has been added as a criteria for CPR</p> <p>Consider advanced airway management if prolonged CPR has been removed.</p>	<p>✓</p> <p>✓</p> <p>✓</p>	<p>x</p> <p>x</p> <p>x</p>
CPG 4.7.31 Allergic Reaction/ Anaphylaxis – Paediatric	<p>Salbutamol NEB is now within the scope of practice for EMTs</p> <p>The conditions for use of Epinephrine auto injector has been changed; it is now indicated for all patients with severe anaphylaxis regardless of whether it has been previously prescribed or not.</p>	<p>✓</p> <p>✓</p>	<p>✓</p> <p>x</p>
CPG 4.7.32 Glycaemic Emergency – Paediatric	<p>A dose of Glucose gel for > 8 year olds has been added</p>	<p>✓</p>	<p>x</p>
CPG 4.7.33 Seizure/ Convulsion – Paediatric	<p>Paracetamol has been removed and replaced with a direction to go to the pyrexia CPG</p>	<p>✓</p>	<p>x</p>

APPENDIX 4 CPG UPDATES FOR EMERGENCY MEDICAL TECHNICIANS

CPGs	The principal differences are	Theory	Skills
CPG 4/5/6.7.50 External Haemorrhage – Paediatric	<p>This CPG has been updated to reflect the importance of managing catastrophic haemorrhage immediately</p> <p>Dressings impregnated with haemostatic agents are now within the scope of practice for EMTs, paramedics and advanced paramedics</p> <p>EMTs, who have completed the BTEC course, may be privileged by a licenced CPG provider to apply a tourniquet</p>	<p>✓</p> <p>✓</p> <p>✓</p>	<p>x</p> <p>✓</p> <p>BTEC only</p>
CPG 4.7.51 Shock from Blood Loss – Paediatric	<p>The entry to this CPG has been changed from 'shock' to 'signs of poor perfusion'</p> <p>An additional care management step has been introduced; Lie patient flat and elevate the legs (if safe to do so)</p>	<p>✓</p> <p>✓</p>	<p>x</p> <p>x</p>
CPG 4.7.52 Spinal Immobilisation – Paediatric	'Consider Paramedic' has been changed to 'Notify a paramedic, advanced paramedic or doctor'	✓	x
CPG 4/5/6.7.53 Burns – Paediatric	Add 'Caution with hypothermia'	✓	x
4/5/6.8.1 Major Emergency – First Practitioners on site	<p>Add 'ambulance loading point'</p> <p>Add 'On site co-ordination centre'</p>	<p>✓</p> <p>✓</p>	<p>x</p> <p>x</p>
4/5/6.8.2 Major Emergency – Operational Control	Add information box 'Controller of Operations may be other than ambulance or fire officers, depending on nature of emergency'	✓	x

APPENDIX 4 CPG UPDATES FOR EMERGENCY MEDICAL TECHNICIANS


New CPGs

New CPGs	The new skills and medications incorporated in the CPG are:	Theory	Skills
CPG 4/5/6.3.4 Asthma – Adult	This CPG outlines the care for a patient with an acute asthma episode	✓	✓
CPG 4/5/6.4.27 Sickle Cell Crisis – Adult	This CPG outlines the care for a patient with a sickle cell crisis.	✓	x
CPG 4/5/6.6.4 Harness Induced Suspension Trauma	This CPG outlines, in particular, the correct posture for patients following harness induced suspension trauma.	✓	x
CPG 4/5/6.6.6 Heat Related Emergency – Adult	This CPG outlines the care for a patient with a heat related emergency.	✓	x
CPG 4/5/6.7.12 Asthma – Paediatric	This CPG outlines the care for a paediatric patient with an acute asthma episode.	✓	✓
CPG 4/5/6.7.35 Pyrexia – Paediatric	This CPG outlines the care for a paediatric patient with a pyrexia episode.	✓	x
CPG 4/5/6.7.36 Sickle Cell Crisis – Paediatric	This CPG outlines the care for a paediatric patient with a sickle cell crisis.	✓	x

APPENDIX 5 PRE-HOSPITAL DEFIBRILLATION POSITION PAPER

Defibrillation is a lifesaving intervention for victims of sudden cardiac arrest (SCA). Defibrillation in isolation is unlikely to reverse SCA unless it is integrated into the chain of survival. The chain of survival should not be regarded as a linear process with each link as a separate entity but once commenced with 'early access' the other links, other than 'post return of spontaneous circulation (ROSC) care', should be operated in parallel subject to the number of people and clinical skills available.

Cardiac arrest management process



ILCOR guidelines 2010 identified that without ongoing CPR, survival with good neurological function from SCA is highly unlikely. Defibrillators in AED mode can take up to 30 seconds between analysing and charging during which time no CPR is typically being performed. The position below is outlined to ensure maximum resuscitation efficiency and safety.

Position

1. Defibrillation mode

- 1.1 Advanced paramedics, and health care professionals whose scope of practice permits, should use defibrillators in manual mode for all age groups.
- 1.2 Paramedics may consider using defibrillators in manual mode for all age groups.
- 1.3 EMTs and responders shall use defibrillators in AED mode for all age groups.

2. Hands off time (time when chest compressions are stopped)

- 2.1 Minimise hands off time, absolute maximum 10 seconds.
- 2.2 Rhythm and/or pulse checks in manual mode should take no more than 5 to 10 seconds and CPR should be recommenced immediately.
- 2.3 When defibrillators are charging CPR should be ongoing and only stopped for the time it takes to press the defibrillation button and recommenced immediately without reference to rhythm or pulse checks.
- 2.4 It is necessary to stop CPR to enable some AEDs to analyse the rhythm. Unfortunately this time frame is not standard with all AEDs. As soon as the analysing phase is completed and the charging phase has begun CPR should be recommenced.

APPENDIX 5 PRE-HOSPITAL DEFIBRILLATION POSITION PAPER

3 Energy

- 3.1 Biphasic defibrillation is the method of choice.
- 3.2 Biphasic truncated exponential (BTE) waveform energy commencing at 150 to 200 joules shall be used.
- 3.3 If unsuccessful the energy on second and subsequent shocks shall be as per manufacturer of defibrillator instructions.
- 3.4 Monophasic defibrillators currently in use, although not as effective as biphasic defibrillators, may continue to be used until they reach the end of their lifespan.

4 Safety

- 4.1 For the short number of seconds while a patient is being defibrillated no person should be in contact with the patient.
- 4.2 The person pressing the defibrillation button is responsible for defibrillation safety.
- 4.3 Defibrillation pads should be used as opposed to defibrillation paddles for pre-hospital defibrillation.

5 Defibrillation pad placement

- 5.1 The right defibrillation pad should be placed mid clavicular directly under the right clavicle.
- 5.2 The left defibrillation pad should be placed mid-axillary with the top border directly under the left nipple.
- 5.3 If a pacemaker or Implantable Cardioverter Defibrillator (ICD) is fitted, defibrillator pads should be placed at least 8 cm away from these devices. This may result in anterior and posterior pad placement which is acceptable.

6 Paediatric defibrillation

- 6.1 Paediatric defibrillation refers to patients less than 8 years of age.
- 6.2 Manual defibrillator energy shall commence and continue with 4 joules/Kg.
- 6.3 AEDs should use paediatric energy attenuator systems.
- 6.4 If a paediatric energy attenuator system is not available an adult AED may be used.
- 6.5 It is extremely unlikely to ever have to defibrillate a child less than 1 year old. Nevertheless, if this were to occur the approach would be the same as for a child over the age of 1. The only likely difference being, the need to place the defibrillation pads anterior and posterior, because of the infant's small size.

7 Implantable Cardioverter Defibrillator (ICD)

- 7.1 If an Implantable Cardioverter Defibrillator (ICD) is fitted in the patient, treat as per CPG. It is safe to touch a patient with an ICD fitted even if it is firing.

8 Cardioversion

- 8.1 Advanced paramedics are authorised to use synchronised cardioversion for unresponsive patients with a tachycardia greater than 150.

Published by:
Pre-Hospital Emergency Care Council
Abbey Moat House, Abbey Street,
Naas, Co Kildare, Ireland.

Phone: + 353 (0)45 882042
Fax: + 353 (0)45 882089

Email: info@phecc.ie
Web: www.phecc.ie

Emergency Medical Technician

